

AD-A014 733

APPROXIMATIONS TO THE DISTRIBUTIONS OF THE LIKELIHOOD
RATIO STATISTICS FOR TESTING CERTAIN STRUCTURES ON
THE COVARIANCE MATRICES OF REAL MULTIVARIATE NORMAL
POPULATIONS

J. C. Lee, et al

Technology, Incorporated

Prepared for:

Aerospace Research Laboratories

June 1975

DISTRIBUTED BY:

NTIS

National Technical Information Service
U. S. DEPARTMENT OF COMMERCE

267074

ARL TR 75-0167



**APPROXIMATIONS TO THE DISTRIBUTIONS
OF THE LIKELIHOOD RATIO STATISTICS FOR
TESTING CERTAIN STRUCTURES ON THE
COVARIANCE MATRICES OF REAL MULTIVARIATE
NORMAL POPULATIONS**

TECHNOLOGY INCORPORATED
3821 COLONEL GLENN HIGHWAY
DAYTON, OHIO 45431

APPLIED MATHEMATICS RESEARCH LABORATORY/ARL

JUNE 1975

INTERIM REPORT



Approved for public release; distribution unlimited

AEROSPACE RESEARCH LABORATORIES/LB
Building 450 — Area B
Wright-Patterson Air Force Base, Ohio 45433

**AIR FORCE SYSTEMS COMMAND
United States Air Force**



ADA014733

NOTICES

When Government drawings, specifications, or other data are used for any purpose other than in connection with a definitely related Government procurement operation, the United States Government thereby incurs no responsibility nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data, is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use, or sell any patented invention that may in any way be related thereto.

Organizations or individuals receiving reports via Aerospace Research Laboratories automatic mailing lists should refer to the ARL number of the report received when corresponding about change of address or cancellation. Such changes should be directed to the specific laboratory originating the report. Do not return this copy; retain or destroy.

Reports are not stocked by the Aerospace Research Laboratories. Copies may be obtained from:


National Technical Information Services
Clearinghouse
Springfield, VA 22161

This technical report has been reviewed and is approved for publication.

FOR THE COMMANDER:

Elizabeth Day

ELIZABETH DAY
Technical Documents
and STINFO Office

AGREES FOR	
NTIS	WASH. SOURCE <input type="checkbox"/>
DDC	DDC SOURCE <input type="checkbox"/>
UNCLASSIFIED	<input type="checkbox"/>
RESTRICTED	<input type="checkbox"/>
BY	
DISTRIBUTION/AVAILABILITY CODES	
Dist.	AVAIL. & W. SPECIAL
	

This report has been reviewed and cleared for open publication and public release by the appropriate Office of Information in accordance with AFR 190-12 and DODD 5230.0. There is no objection to unlimited distribution of this report to the public at large, or by DDC to the National Technical Information Service.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER ARL 75-0167	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) APPROXIMATIONS TO THE DISTRIBUTIONS OF THE LIKELIHOOD RATIO STATISTICS FOR TESTING CERTAIN STRUCTURES ON THE COVARIANCE MATRICES OF REAL MULTIVARIATE NORMAL POPULATIONS		5. TYPE OF REPORT & PERIOD COVERED Technical - Interim
7. AUTHOR(s) J. C. Lee T. C. Chang P. R. Krishnaiah		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Technology Incorporated 3821 Colonel Glenn Highway Dayton, Ohio 45431		8. CONTRACT OR GRANT NUMBER(s) F33615-73-C-4155 and In-House Research
11. CONTROLLING OFFICE NAME AND ADDRESS Applied Mathematics Research Lab/LB Aerospace Research Laboratories (AFSC) Wright-Patterson AFB, Ohio 45433		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS DoD Element 61102F 70710212
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE June 1975
		13. NUMBER OF PAGES 54
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) multivariate analysis Pearson type approximations likelihood ratio statistics covariance structures tables		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In this report, the authors approximated the distributions of the likelihood ratio statistics for testing certain hypotheses on the covariance matrices of the multivariate normal populations by using suitable Pearson type distributions. The hypotheses that are considered include the hypotheses of sphericity, multiple homogeneity of covariance matrices and multiple independence. Using these approximations, tables of percentage points of the distributions considered are constructed. The accuracy of these tables is sufficient for practical purposes.		

DD FORM 1473
1 JAN 73

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

PREFACE

This report was prepared for the Applied Mathematics Research Laboratory, Aerospace Research Laboratories, by J. C. Lee, T. C. Chang and P. R. Krishnaiah under Project 7071, "Research in Applied Mathematics", Work Unit 12, "Multivariate Analysis and Its Applications". The work of J. C. Lee and T. C. Chang was performed at the Aerospace Research Laboratories in the capacity of Technology Incorporated Visiting Research Associates under Contract F33615-73-C-4155. J. C. Lee and T. C. Chang are currently on leave of absence from Wright State University and University of Cincinnati, respectively.

TABLE OF CONTENTS

SECTION		PAGE
I	INTRODUCTION	1
II	STATEMENT OF THE PROBLEMS AND PRELIMINARIES	3
III	MULTIPLE INDEPENDENCE OF SEVERAL SETS OF VARIABLES	5
IV	LIKELIHOOD RATIO TEST STATISTIC FOR SPHERICITY	14
V	LIKELIHOOD RATIO STATISTIC FOR TESTING THE HYPOTHESIS THAT THE COVARIANCE MATRIX IS EQUAL TO A SPECIFIED MATRIX	16
VI	LIKELIHOOD RATIO STATISTIC FOR TESTING THE HOMOGENEITY OF COVARIANCE MATRICES	19
VII	LIKELIHOOD RATIO TEST STATISTIC FOR MULTIPLE HOMOGENEITY OF COVARIANCE MATRICES	30
	REFERENCES	37
	APPENDIX	39

LIST OF TABLES

TABLE		PAGE
I	Comparison of the Pearson Type Approximation with the Asymptotic Expansion of Order $n^{-1/3}$ for the Distribution of V_1^*	9
II	Comparison of the Pearson Type Approximation with Inversion of the Asymptotic Expression for the Percentage Points of V_1^*	10
III	Percentage Points of $-2 \log V_1$	11
IV	Comparison of the Pearson Type Approximation with Exact Expression for the Distribution of V_2	15
V	Comparison of the Pearson Type Approximation with Exact and Asymptotic Expressions for the Distribution of V_3^*	18
VI	Comparison of the Pearson Type Approximation with the Asymptotic Expression for the Distribution of V_4^*	21
VII	Percentage Points of the Distribution of $-2 \log V_4$	22
VIII	Comparison of the Pearson Type Approximation with the Asymptotic Expression for the Distribution of V_5^*	32
IX	Percentage Points of V_5^*	33

SECTION I

INTRODUCTION

The problems of testing the hypotheses on the structures of the covariance matrix of the multivariate normal population have received considerable attention in the literature since these problems have applications in various disciplines. Wilks [1,2] is one of the earliest workers in this area. In this report, we investigate approximations to the distributions of the likelihood ratio statistics for testing the hypotheses of (1) multiple independence of several sets of variables, (2) sphericity, (3) equality of the covariance matrix to a given matrix, and (4) equality of the covariance matrices of independent sets of variables. These approximations are based upon fitting suitable Pearson type distributions by using the first four moments of the above statistics. Using the above approximations, we construct percentage points of these statistics. The accuracy of these approximations is sufficient for practical purposes.

Exact distributions of the test statistics considered here are very difficult to compute. One can, of course, use Box's asymptotic expression by taking a sufficient number of terms to compute the tables and this may be preferable to exact expressions from a computational point of view. However, this asymptotic expression is also difficult to compute when we need to take several terms in the series and this is the case when the sample size is not large. Since Pearson type approximations are quite simple from a computational point of view and since their accuracy is sufficient for practical purposes, they are definitely preferable to either exact expressions or Box's asymptotic series if we

are interested in computing the percentage points of the statistics considered in this paper.

SECTION II

STATEMENT OF THE PROBLEMS AND PRELIMINARIES

Let $\underline{X}' = (\underline{X}'_1, \dots, \underline{X}'_q)$ be distributed as a multivariate normal with mean vector $\underline{\mu}'$ and covariance matrix Σ . Also, let $E(\underline{X}_i) = \underline{\mu}_i$ and $E\{(\underline{X}_i - \underline{\mu}_i)(\underline{X}_j - \underline{\mu}_j)'\} = \Sigma_{ij}$, where \underline{X}_i is of order $p_i \times 1$ and $s = \sum_{i=1}^q p_i$.

In this paper, we consider approximations to the distributions of the likelihood ratio statistics for testing the hypotheses H_1, H_2, H_3, H_4 , and H_5 where

$$H_1: \Sigma_{ij} = 0 \quad (i \neq j = 1, \dots, q)$$

$$H_2: \Sigma = \sigma^2 \Sigma_0 \quad (\sigma^2 \text{ is unknown, } \Sigma_0 \text{ is known})$$

$$H_3: \Sigma = \Sigma_0$$

$$H_4: \Sigma_{11} = \dots = \Sigma_{qq} \quad (\text{under the assumption that } H_1 \text{ is true})$$

$$\text{and } p_1 = \dots = p_q$$

$$H_5: \left\{ \begin{array}{l} \Sigma_{11} = \dots = \Sigma_{q_1, q_1} \\ \Sigma_{q_1+1, q_1+1} = \dots = \Sigma_{q_2^*, q_2^*} \\ \Sigma_{q_{k-1}^*+1, q_{k-1}^*+1} = \dots = \Sigma_{q, q} \end{array} \right.$$

Here $q_0^* = 0$, $q_j^* = \sum_{i=1}^j q_i$ and $q_k^* = q$.

These distributions are approximated with Pearson type distributions. A brief description of the family of Pearson type distributions is given below.

If $f(x)$, $(c < x < d)$, is the probability density function belonging to the family of Pearson type distributions, then $f(x)$ satisfies the differential equation

$$(b_0 + b_1x + b_2x^2) \frac{df(x)}{dx} = (x + a)f(x). \quad (1)$$

Let $m'_k = \int_c^d x^k f(x) dx$ and $m_k = \int_c^d (x - m'_1)^k f(x) dx$. It is known that

$$\begin{aligned} a &= m_3(m_4 + 3m_2^2)/A \\ b_0 &= -m_2(4m_2m_4 - 3m_3^2)/A \\ b_1 &= -m_3(m_4 + 3m_2^2)/A \\ b_2 &= -(2m_2m_4 - 3m_3^2 - 6m_2^3)/A \\ A &= 10m_2m_4 - 18m_2^3 - 12m_3^2 \end{aligned} \quad (2)$$

The family of Pearson type distributions is completely specified by its first four moments and the type of the distribution can be determined by examining the quantity κ where

$$\kappa = \frac{\beta_1(\beta_2 + 3)^2}{4(2\beta_2 - 3\beta_1 - 6)(4\beta_2 - 3\beta_1)}, \quad (3)$$

$\beta_1 = m_3^2/m_2^3$ and $\beta_2 = m_4/m_2^2$. If κ is negative, we classify it as the Pearson's Type I distribution. When κ is positive, it is classified as the Pearson's Type IV or Type VI accordingly as κ is less than or greater than unity, respectively. For a discussion of Pearson type distributions, see [3]. Tables for the percentage points of Pearson type distributions are given in Johnson, Nixon and Amos [4] for some values of the parameters.

SECTION III
MULTIPLE INDEPENDENCE OF SEVERAL
SETS OF VARIABLES

Let $\tilde{x}_j' = (\tilde{x}_{1j}', \dots, \tilde{x}_{qj}')$, $(j=1, \dots, N)$, be j th independent observation on \tilde{x}' and let

$$A = \begin{bmatrix} A_{11} & \dots & A_{1q} \\ A_{21} & \dots & A_{2q} \\ \cdot & \dots & \cdot \\ \cdot & \dots & \cdot \\ \cdot & \dots & \cdot \\ A_{q1} & \dots & A_{qq} \end{bmatrix}$$

where

$$A_{lm} = \sum_{j=1}^N (\tilde{x}_{lj} - \bar{\tilde{x}}_{l\cdot}) (\tilde{x}_{mj} - \bar{\tilde{x}}_{m\cdot})'$$

and $\bar{\tilde{x}}_{l\cdot} = \frac{1}{N} \sum_{j=1}^N \tilde{x}_{lj}$. Then, the likelihood ratio statistic for testing

H_1 is known to be

$$V_1 = \frac{|A|}{\prod_{j=1}^q |\Lambda_{jj}|} \quad (4)$$

The moments of the statistic V_1 are given by

$$E(V_1^h) = \frac{\prod_{i=1}^s \Gamma \left[\frac{1}{2}(n+1-i) + h \right] \prod_{i=1}^q \left\{ \prod_{j=1}^{p_i} \Gamma \left[\frac{1}{2}(n+1-j) \right] \right\}}{\prod_{i=1}^s \Gamma \left[\frac{1}{2}(n+1-i) \right] \prod_{i=1}^q \left\{ \prod_{j=1}^{p_i} \Gamma \left[\frac{1}{2}(n+1-j) + h \right] \right\}} \quad (5)$$

where $n = N-1$ and $\Gamma(n)$ is the complete gamma function. The statistic V_1 and its moments were derived by Wilks [2]. According to the likelihood ratio test, we accept or reject H_1 accordingly as $V_1^* \lessgtr c$, where $V_1^* = -2\log V_1$ and

$$P[V_1^* \leq c_1 | H_1] = (1 - \alpha) \quad (6)$$

The distribution of V_1 is quite skew. Hence, we approximate the distribution of $V_1^{1/b}$ with a suitable Pearson type distribution, where b is a properly chosen integer. The type of Pearson distribution that has to be fitted is determined by computing κ given by Eq. (3) after replacing m'_1, m'_2, m'_3 , and m'_4 with $\mu'_{b1}, \mu'_{b2}, \mu'_{b3}$ and μ'_{b4} , respectively, where μ'_{bh} is the h -th moment of $V_1^{1/b}$. By the above method, it is found that the Pearson's Type I curve fits the distribution of $V_1^{1/b}$. The density of the Pearson's Type I distribution is given by

$$f(x) = \frac{1}{\beta(\alpha+1, \epsilon+1)(\sigma-\sigma_0)^{\alpha+\epsilon+1}} (x-\sigma_0)^\alpha (\sigma-x)^\epsilon \quad (7)$$

$$\sigma_0 \leq x \leq \sigma$$

where σ_0, α, σ and ϵ depend upon a^0, b_0^0, b_1^0 and b_2^0 . The parameters a^0, b_0^0, b_1^0 and b_2^0 are respectively equivalent to a, b_0, b_1 , and b_2 after replacing m'_i with μ'_{bi} . In this report we consider the case where $p_i = p$.

Box [5] derived asymptotic expressions for a class of likelihood ratio test statistics in multivariate statistical analysis. This class includes the statistics associated with testing the hypotheses H_1, H_2, H_4 , and H_5 . The number of terms given by Box is not sufficient to get

the desired degree of accuracy in several practical situations. So, the authors gave terms up to $O(n^{-15})$ in the appendix applying the method of Box.

Consul [6] and Mathai and Rathie [7] derived exact expressions for the distribution of V_1 , and these expressions are very difficult to compute. So, the authors studied approximations to these distributions with suitable Pearson type distributions.

Table I gives a comparison of values obtained by the Pearson type approximation and the asymptotic expression of order n^{-13} . In Table I, α_1 is the value of α if we use the Pearson type approximation whereas α_2 is the value of α when we use the asymptotic expression of order n^{-13} , where α is given by

$$P[V_1^* \leq c_1 | H_1] = (1-\alpha). \quad (8)$$

Davis and Field [8] computed the percentage points of $-2p \log V_1$ for some values of the parameters by using the Cornish-Fisher type inversion (see Davis [9]) of Box's asymptotic series when

$$1-\rho = \left\{ 2 \left(s^3 - \sum_{i=1}^q p_i^3 \right) + 9 \left(s^2 - \sum_{i=1}^q p_i^2 \right) \right\} / 6N \left(s^2 - \sum_{i=1}^q p_i^2 \right)$$

In Table II, the entries under the columns L-C-K are the values of c_1 obtained by the authors with the Pearson type approximation whereas the entries under the columns D-F are the corresponding values obtained by Davis and Field [8].

Tables I and II indicate that the accuracy of the Pearson type approximation to the distribution of V_1 is sufficient for practical purposes. Hence, using the Pearson type approximation, we computed the values of c_1 for $p = 1, 2, 3$; $q = 3, 4, 5$; $\alpha = 0.01, 0.05, 0.10$; and $M = 1(1)20(2)30$, where $M = n - s - 3$. These values are given in Table III. When $q = 2$, it is found that the Pearson type approximation is quite satisfactory, and the results are reported in a companion paper [10] by the authors.

TABLE I
COMPARISON OF THE PEARSON TYPE APPROXIMATION
WITH THE ASYMPTOTIC EXPANSION OF ORDER $n^{-1/3}$
FOR THE DISTRIBUTION OF V_1^*

n	q=3 p=1			q=5 p=1		
	c_1	α_1	α_2	c_1	α_1	α_2
10	1.913	0.05	0.0499	4.978	0.05	0.0488
10	2.778	0.01	0.0100	6.333	0.01	0.0095
15	1.187	0.05	0.0500	2.947	0.05	0.0497
15	1.723	0.01	0.0100	3.742	0.01	0.0099
20	0.860	0.05	0.0500	2.099	0.05	0.0499
20	1.249	0.01	0.0100	2.663	0.01	0.0100
30	0.555	0.05	0.0500	1.333	0.05	0.0500
30	0.805	0.01	0.0100	1.690	0.01	0.0100

TABLE II
COMPARISON OF THE PEARSON TYPE APPROXIMATION WITH
INVERSION OF THE ASYMPTOTIC EXPRESSION FOR
THE PERCENTAGE POINTS OF V_1^*

		q=4	p=1	q=5	p=1
n	α	L-C-K	D-F	L-C-K	D-F
10	0.05	3.238	3.238	4.978	4.977
10	0.01	4.330	4.331	6.333	6.333
15	0.05	1.967	1.967	2.949	2.948
15	0.01	2.628	2.628	3.742	3.742
20	0.05	1.414	1.414	2.099	2.099
20	0.01	1.888	1.888	2.663	2.663
24	0.05	1.154	1.154	1.706	1.707
24	0.01	1.541	1.542	2.165	2.164

TABLE III*
PERCENTAGE POINTS OF $-2 \log V_1$
q=3

M	p=1			p=2			p=3		
	α	10%	5%	1%	10%	5%	1%	10%	5%
1		2.419	3.023	4.390	5.686	6.371	7.991	9.161	10.037
2		2.027	2.534	3.670	4.851	5.509	6.899	8.091	8.857
3		1.744	2.100	3.165	4.279	4.857	6.076	7.255	7.937
4		1.531	1.913	2.770	3.829	4.345	5.432	6.502	7.190
5		1.363	1.704	2.475	3.466	3.933	4.914	6.026	6.587
6		1.229	1.537	2.231	3.167	3.593	4.487	5.559	6.075
7		1.119	1.400	2.031	2.916	3.307	4.129	5.160	5.639
8		1.027	1.284	1.864	2.701	3.064	3.825	4.815	5.262
9		.949	1.167	1.723	2.517	2.854	3.563	4.515	4.933
10		.882	1.103	1.682	2.356	2.672	3.334	4.250	4.643
11		.824	1.030	1.496	2.215	2.511	3.134	4.016	4.386
12		.773	.967	1.403	2.089	2.369	2.956	3.805	4.157
13		.728	.910	1.322	1.977	2.242	2.798	3.616	3.950
14		.688	.850	1.249	1.877	2.120	2.655	3.445	3.763
15		.652	.815	1.183	1.786	2.025	2.527	3.289	3.593
16		.620	.775	1.125	1.704	1.932	2.410	3.148	3.434
17		.591	.739	1.072	1.629	1.847	2.304	3.018	3.296
18		.564	.705	1.024	1.561	1.769	2.207	2.898	3.165
19		.540	.675	.979	1.497	1.697	2.117	2.788	3.044
20		.517	.647	.939	1.439	1.631	2.035	2.685	2.932
22		.478	.597	.867	1.335	1.513	1.880	2.501	2.732
24		.446	.555	.805	1.245	1.412	1.761	2.341	2.557
26		.414	.510	.752	1.167	1.323	1.650	2.200	2.403
28		.388	.486	.705	1.098	1.244	1.552	2.076	2.267
30		.366	.458	.664	1.036	1.175	1.465	1.965	2.145

* The entries in this table are the values of c_1 where $M = n - s - 3$ and $p[-2 \log V_1 \leq c_1] = (1 - \alpha)$.

TABLE Iii (Continued)

q = 4

M	α	p=1			p=2			p=3		
		10%	5%	1%	10%	5%	1%	10%	5%	1%
1		3.694	4.376	5.061	0.557	9.417	11.195	13.949	14.932	16.905
2		3.143	3.721	4.980	7.534	8.285	9.831	12.539	13.408	15.155
3		2.736	3.238	4.330	6.739	7.406	8.777	11.407	12.190	13.762
4		2.422	2.867	3.832	6.100	6.701	7.936	10.473	11.189	12.618
5		2.174	2.572	3.438	5.574	6.122	7.245	9.609	10.347	11.660
6		1.971	2.333	3.118	5.134	5.637	6.669	9.017	9.627	10.846
7		1.804	2.134	2.852	4.769	5.225	6.179	8.436	9.004	10.140
8		1.663	1.967	2.628	4.436	4.870	5.757	7.927	8.460	9.524
9		1.542	1.825	2.437	4.154	4.560	5.391	7.477	7.980	8.981
10		1.438	1.701	2.272	3.907	4.289	5.068	7.078	7.552	8.498
11		1.347	1.593	2.128	3.688	4.048	4.783	6.719	7.169	8.066
12		1.266	1.498	2.001	3.493	3.833	4.528	6.395	6.823	7.676
13		1.195	1.414	1.888	3.317	3.639	4.299	6.103	6.510	7.323
14		1.132	1.338	1.788	3.158	3.465	4.093	5.835	6.225	7.001
15		1.074	1.271	1.697	3.013	3.306	3.905	5.591	5.964	6.708
16		1.023	1.210	1.616	2.882	3.162	3.734	5.367	5.725	6.437
17		.976	1.154	1.541	2.761	3.030	3.578	5.160	5.503	6.188
18		.933	1.104	1.474	2.650	2.908	3.434	4.968	5.299	5.958
19		.894	1.057	1.412	2.548	2.796	3.302	4.790	5.110	5.745
20		.858	1.015	1.355	2.453	2.692	3.179	4.626	4.933	5.546
22		.794	.939	1.254	2.284	2.506	2.959	4.327	4.615	5.188
24		.739	.874	1.167	2.137	2.344	2.768	4.066	4.336	4.874
26		.691	.817	1.091	2.007	2.202	2.600	3.833	4.088	4.596
28		.649	.767	1.025	1.893	2.076	2.451	3.627	3.868	4.348
30		.612	.723	.966	1.790	1.964	2.318	3.442	3.671	4.125

TABLE III (Continued)

q = 5

M	p=1			p=2			p=3		
	α	10X	5X	1X	10X	5X	1X	10X	5X
1	5.039	5.784	7.368	11.665	12.600	14.500	16.945	20.006	22.106
2	4.339	4.978	6.333	10.405	11.228	12.899	17.242	18.192	20.067
3	3.613	4.371	5.557	9.405	10.146	11.639	15.852	16.716	18.421
4	3.402	3.908	4.954	8.589	9.261	10.616	14.687	15.477	17.043
5	3.071	3.520	4.478	7.909	8.525	9.767	13.691	14.425	15.876
6	2.801	3.208	4.073	7.331	7.900	9.047	12.828	13.515	14.864
7	2.573	2.949	3.742	6.835	7.364	8.430	12.073	12.717	13.981
8	2.381	2.727	3.461	6.402	6.897	7.893	11.406	12.011	13.204
9	2.215	2.537	3.219	6.023	6.487	7.423	10.811	11.386	12.513
10	2.071	2.372	3.009	5.686	6.125	7.006	10.279	10.821	11.890
11	1.944	2.227	2.826	5.386	5.801	6.635	9.798	10.313	11.332
12	1.832	2.099	2.663	5.116	5.510	6.301	9.360	9.852	10.822
13	1.733	1.985	2.518	4.873	5.247	6.000	8.960	9.432	10.359
14	1.644	1.882	2.388	4.652	5.009	5.727	8.594	9.047	9.935
15	1.563	1.790	2.270	4.449	4.791	5.477	8.259	8.691	9.545
16	1.490	1.706	2.165	4.265	4.592	5.249	7.948	8.364	9.185
17	1.424	1.631	2.068	4.095	4.409	5.039	7.661	8.061	8.858
18	1.363	1.561	1.988	3.938	4.240	4.846	7.393	7.779	8.541
19	1.307	1.497	1.898	3.793	4.084	4.667	7.144	7.518	8.252
20	1.256	1.438	1.823	3.658	3.938	4.501	6.911	7.272	7.983
22	1.164	1.333	1.698	3.415	3.677	4.282	6.498	6.828	7.495
24	1.085	1.243	1.576	3.203	3.448	3.948	6.116	6.435	7.063
26	1.016	1.163	1.475	3.016	3.247	3.789	5.784	6.086	6.679
28	.955	1.094	1.387	2.849	3.067	3.584	5.486	5.772	6.335
30	.901	1.032	1.309	2.788	2.986	3.321	5.219	5.491	6.025

SECTION IV

LIKELIHOOD RATIO TEST STATISTIC FOR SPHERICITY

The likelihood ratio statistic for testing $H_2 : \Sigma = \sigma^2 \Sigma_0$, where σ^2 is unknown and Σ_0 is known, is given by

$$V_2 = \frac{|A \Sigma_0^{-1}|}{\left\{ \text{tr } A \Sigma_0^{-1} / s \right\}^s} \quad (9)$$

where A was defined in Section III and $\text{tr } B$ denotes the trace of B . The h -th moment of V_2 is given by

$$E(V_2^h) = \frac{p^{hs} \Gamma(sn/2)}{\Gamma(sh + \frac{sn}{2})} \prod_{i=1}^s \frac{\Gamma\left[\frac{1}{2}(n+1-i)+h\right]}{\Gamma\left[\frac{1}{2}(n+1-i)\right]} \quad (10)$$

The statistic V_2 and its moments were derived by Mauchly [11]. Using the first four moments, we approximated the distribution of $V_2^{1/b}$ with the Pearson Type I distribution, where b is a suitably chosen integer. Expressions for the exact distribution of V_2 were given by Consul [6], Mathai and Rathie [12], and Nagarsenker and Pillai [13], but these expressions are very difficult to compute. In Table IV the values under the column LCK are the values obtained by the authors using the Pearson type approximation whereas the corresponding exact values are taken from Nagarsenker and Pillai [13]. This table indicates that the accuracy of the Pearson type approximation is sufficient for practical purposes. The values of α in this table are given by the relation

$$P \left[V_2 \geq c_2 | H_2 \right] = (1 - \alpha)$$

TABLE IV
COMPARISON OF THE PEARSON TYPE APPROXIMATION WITH
EXACT EXPRESSION FOR THE DISTRIBUTION OF V_2

n	α	s=4		s=5		s=7	
		L-C-K	Exact	L-C-K	Exact	L-C-K	Exact
6	0.05	0.0169	0.0169	0.0013	0.0013	--	--
6	0.01	0.0050	0.0050	0.00022	0.00022	--	--
10	0.05	0.1297	0.1297	0.0492	0.0492	0.0029	0.0030
10	0.01	0.0726	0.0726	0.0242	0.0242	0.0010	0.0010
15	0.05	0.2812	0.2812	0.1608	0.1608	0.0368	0.0368
15	0.01	0.1967	0.1966	0.1052	0.1052	0.0207	0.0207
21	0.05	0.4173	0.4173	0.2877	0.2876	0.1111	0.1111
21	0.01	0.3264	0.3264	0.2156	0.2156	0.0761	0.0761
33	0.05	0.5833	0.5833	0.4663	0.4663	0.2665	0.2665
33	0.01	0.5013	0.5013	0.3910	0.3910	0.2125	0.2125
41	0.05	0.6507	0.6508	0.5453	0.5453	0.3515	0.3515
41	0.01	0.5769	0.5769	0.4741	0.4741	0.2939	0.2939

SECTION V

LIKELIHOOD RATIO STATISTIC FOR TESTING THE
HYPOTHESIS THAT THE COVARIANCE MATRIX
IS EQUAL TO A SPECIFIED MATRIX

Consider the problem of testing the hypothesis $H_3 : \Sigma = \Sigma_0$ where Σ_0 is specified. The likelihood ratio statistic for testing H_3 and the moments of this statistic were derived by Anderson [14]. The modified likelihood ratio test statistic (obtained by changing N to n in the likelihood ratio statistic) and its moments are as given below:

$$V_3 = (e/n)^{sn/2} \left| A \Sigma_0^{-1} \right|^{n/2} \text{etr} \left(-\frac{1}{2} A \Sigma_0^{-1} \right) \quad (11)$$

$$E(V_3^h) = (2e/n)^{shn/2} \left| \Sigma_0 \right|^{nh/2} \cdot \left| I + h \Sigma_0 \right|^{-n(1+h)/2} \frac{\prod_{i=1}^s \Gamma \left[\frac{1}{2}(n+nh+1-i) \right]}{\prod_{i=1}^s \Gamma \left[\frac{1}{2}(n+1-i) \right]} \quad (12)$$

Using the first four moments, we approximated the distribution of $V_3^{1/b}$ with the Pearson Type I distribution as in Section III, where b is a suitably chosen integer. Korin [15] obtained an asymptotic expression of order n^{-15} for the distribution of $V_3^* = -2 \log V_3$. Using this expression, he computed percentage points of V_3^* for some values of the parameters. Nagarsenker and Pillai [16] obtained an expression for the distribution of V_3^* , but this expression is complicated from a computational point of view. Using this expression, they computed exact percentage points of V_3^* . In Table V we compare our values (given in the column L-C-K) obtained by using the Pearson type approximation with exact

values of Nagarsenker and Pillai, and the values obtained by Korin [15]. Table V indicates that the accuracy of the Pearson type approximation is sufficient for practical purposes.

TABLE V
COMPARISON OF THE PEARSON TYPE APPROXIMATION
WITH EXACT AND ASYMPTOTIC EXPRESSIONS
FOR THE DISTRIBUTION OF V_3^*

s=4					s=6					s=10				
n	α	L-C-K	Exact	Korin	n	α	L-C-K	Exact	Korin	n	α	L-C-K	Exact	
6	0.05	25.76	25.76	25.8	8	0.05	49.24	49.25	--	12	0.05	119.08	119.07	
6	0.01	33.07	33.08	--	8	0.01	59.59	59.60	--	12	0.01	135.81	135.81	
7	0.05	24.06	24.06	24.06	9	0.05	45.82	45.83	--	13	0.05	111.15	111.15	
7	0.01	30.74	30.75	30.8	9	0.01	55.15	55.16	--	13	0.01	126.10	126.10	
10	0.05	21.75	21.75	21.75	10	0.05	43.62	43.63	--	14	0.05	105.76	105.76	
10	0.01	27.66	27.66	27.66	10	0.01	52.35	52.35	--	14	0.01	119.64	119.64	
11	0.05	21.35	21.35	21.35	15	0.05	39.71	38.71	--	15	0.05	101.83	101.82	
11	0.01	27.13	27.13	27.13	15	0.01	46.23	46.23	--	15	0.01	114.96	114.97	
13	0.05	20.77	20.77	20.77	20	0.05	36.87	36.86	36.87	20	0.05	91.28	91.28	
13	0.01	26.38	--	--	20	0.01	43.97	43.98	43.99	20	0.01	102.71	102.70	
					25	0.05	35.89	35.88	35.89	25	0.05	86.51	86.52	
					25	0.01	42.79	42.79	42.80	25	0.01	97.23	97.23	

SECTION VI
 LIKELIHOOD RATIO STATISTIC FOR TESTING
 THE HOMOGENEITY OF COVARIANCE MATRICES

Let $\tilde{X}' = (\tilde{X}'_1, \dots, \tilde{X}'_q)$ be as defined in Section II. Let \tilde{X}_{ij} ($j=1, \dots, N_i$) be j -th independent observation on \tilde{X}_i . Also, let $p_i = p(i = 1, \dots, q)$ and

$$A_{ii} = \sum_{j=1}^{N_i} (\tilde{X}_{ij} - \bar{\tilde{X}}_i)(\tilde{X}_{ij} - \bar{\tilde{X}}_i)'$$

Wilks [1] derived the likelihood ratio statistic for $H_4 : \Sigma_{11} = \dots = \Sigma_{qq}$ when $\Sigma_{ij} = 0 (i \neq j=1, \dots, q)$ and derived its moments. Let $n_i = N_i - 1$, and $n = \sum_{i=1}^q n_i$. The modified likelihood ratio statistic V_4 (obtained by interchanging N_i with n_i in the likelihood ratio statistic) for H_4 and the moments of V_4 are given below:

$$V_4 = \frac{\prod_{g=1}^q |A_{gg}|^{n_g/2}}{\left| \prod_{g=1}^q A_{gg} \right|^{n/2}} \frac{n^{pn/2}}{\prod_{g=1}^q n_g^{pn_g/2}} \quad (13)$$

$$E(V_4^h) = \left(n^{phn/2} / \prod_{g=1}^q n_g^{phn_g/2} \right)$$

$$\begin{aligned} & \cdot \prod_{i=1}^p \left[\left\{ \prod_{g=1}^q \frac{\Gamma((n_g + hn_g + 1 - i)/2)}{\Gamma((n_g + 1 - i)/2)} \right\} \right. \\ & \cdot \left. \left\{ \frac{\Gamma((n + 1 - i)/2)}{\Gamma((n + hn + 1 - i)/2)} \right\} \right] \end{aligned} \quad (14)$$

Korin [17] computed percentage points by using Box's asymptotic expression up to terms of order n^{-15} . Davis and Field [8] computed the percentage points by using the Cornish-Fisher type inversion of Box's asymptotic expression. In this paper, we approximated the distribution of $V_4^{1/b}$, with a suitable Pearson type distribution where b is a suitably chosen integer. The type of distribution that is fitted is the Pearson's Type I distribution. In Table VI, we compared some of the percentage points obtained by us with the corresponding values obtained by Davis and Field [8] and Korin [17], where $n_i = n_0$. The percentage points c_4 obtained by us using the Pearson type approximation are given under the column L-C-K whereas the corresponding values obtained by Korin, and Davis-Field are given under the columns Korin and D-F, respectively. Table VI indicates that the Pearson type approximation is satisfactory for practical purposes. In this table, α is defined by the equation

$$P \left[V_4^* \leq c_4 | H_4 \right] = (1 - \alpha) \quad (15)$$

When n_0 is large, Bishop [18] investigated the accuracy of approximating the distribution of the $\frac{2}{N}$ th power of the likelihood ratio criterion with the Beta distribution for a few cases, where $n = qn_0$.

Using the Pearson type approximation, we computed the values of c_4 when $n_0 = (p+1)(1)20(5)30$, $p = 2(1)6$, $q = 2(1)10$ and $\alpha = 0.01, 0.05, 0.10$. These percentage points are given in Table VII.

TABLE VI
COMPARISON OF THE PEARSON TYPE APPROXIMATION
WITH THE ASYMPTOTIC EXPRESSION FOR
THE DISTRIBUTION OF V_4^*

$\alpha \approx 0.05$

n_0	k	$p=2$		$p=3$		$p=6$	
		L-C-K	Korin	L-C-K	Korin	L-C-K	D-F
			D-F		D-F		D-F
4	2	10.70	10.70	22.41	--	--	--
4	9	46.07	46.07	99.94	--	--	--
7	2	9.24	9.24	16.59	16.59	69.62	--
7	9	41.26	41.26	79.90	79.91	345.81	--
10	2	8.76	8.76	15.11	15.11	49.95	49.95
10	9	39.65	36.65*	74.58	74.57	271.84	--
15	2	8.42	--	14.15	--	42.03	42.02
15	9	38.50	--	71.05	--	239.72	--
20	2	8.26	--	13.72	--	39.11	39.10
20	9	37.95	--	69.45	--	227.45	--

* There is a typographical error in Korin's table. The correct value seems to be 39.65.

TABLE VII
PERCENTAGE POINTS OF THE DISTRIBUTION
OF $-2 \log V_4$

$\alpha = 0.10$ $p = 2$

n_0	$q=2$	$q=3$	$q=4$	$q=5$	$q=6$	$q=7$	$q=8$	$q=9$	$q=10$
3	9.75	15.30	21.29	26.52	31.50	36.58	41.43	46.32	51.12
4	8.56	14.07	19.08	23.87	28.52	33.07	37.55	41.97	46.36
5	7.97	13.22	17.99	22.55	26.37	31.31	35.58	39.80	43.97
6	7.63	12.70	17.33	21.75	26.05	30.25	34.39	38.49	42.53
7	7.39	12.36	16.89	21.22	25.42	29.54	33.60	37.61	41.57
8	7.23	12.11	16.57	20.84	24.78	29.04	33.03	36.98	40.88
9	7.10	11.93	16.34	20.55	24.35	28.66	32.61	36.51	40.37
10	7.01	11.78	16.16	20.33	24.19	28.36	32.28	36.14	39.97
11	6.93	11.67	16.01	20.15	24.18	28.12	32.01	35.85	39.65
12	6.87	11.58	15.88	20.01	24.01	27.93	31.79	35.61	39.38
13	6.82	11.50	15.79	19.89	23.87	27.77	31.62	35.41	39.16
14	6.77	11.43	15.70	19.78	23.75	27.63	31.46	35.24	38.98
15	6.74	11.37	15.63	19.69	23.65	27.52	31.33	35.09	38.82
16	6.71	11.33	15.57	19.62	23.56	27.42	31.21	34.97	38.68
17	6.68	11.28	15.51	19.55	23.48	27.32	31.11	34.86	38.56
18	6.65	11.25	15.46	19.49	23.41	27.25	31.03	34.70	38.45
19	6.63	11.21	15.42	19.44	23.35	27.18	30.95	34.68	38.36
20	6.61	11.18	15.38	19.40	23.30	27.12	30.88	34.60	38.28
25	6.53	11.07	15.24	19.22	23.09	26.88	30.61	34.30	37.95
30	6.48	11.00	15.14	19.10	22.96	26.73	30.45	34.12	37.74

$\alpha = 0.10$ $p = 3$

n_0	$q=2$	$q=3$	$q=4$	$q=5$	$q=6$	$q=7$	$q=8$	$q=9$	$q=10$
4	18.87	30.76	41.78	52.41	62.90	73.03	83.14	93.16	103.12
5	16.19	26.87	36.79	46.36	55.73	64.95	74.05	83.09	92.05
6	14.83	24.87	34.22	43.24	52.06	60.76	69.35	77.86	86.30
7	14.01	23.66	32.64	41.33	49.91	58.18	66.44	74.64	82.76
8	13.46	22.83	31.56	40.03	48.29	56.44	64.48	72.45	80.37
9	13.06	22.24	30.81	39.09	47.19	55.17	63.05	70.88	78.64
10	12.77	21.79	30.23	38.38	46.36	54.21	61.93	69.68	77.31
11	12.53	21.44	29.77	37.82	45.70	53.46	61.13	68.74	76.28
12	12.35	21.16	29.40	37.37	45.17	52.86	60.45	67.98	75.45
13	12.19	20.93	29.10	37.00	44.74	52.36	59.83	67.35	74.77
14	12.07	20.73	28.85	36.69	44.38	51.94	59.42	66.84	74.19
15	11.96	20.57	28.63	36.43	44.16	51.59	59.03	66.39	73.70
16	11.87	20.43	28.45	36.21	43.90	51.28	58.67	66.01	73.29
17	11.78	20.30	28.29	36.01	43.57	51.02	58.38	65.67	72.92
18	11.71	20.20	28.15	35.83	43.37	50.79	58.11	65.38	72.59
19	11.65	20.10	28.02	35.68	43.19	50.58	57.98	65.12	72.32
20	11.60	20.02	27.91	35.55	43.02	50.39	57.68	64.90	72.07
25	11.39	19.70	27.50	35.05	42.43	49.71	56.91	64.04	71.12
30	11.26	19.50	27.23	34.72	42.05	49.27	56.42	63.43	70.52

TABLE VII (Continued)

 $\alpha = 0.10 \quad p = 4$

n_o	q=2	q=3	q=4	q=5	q=6	q=7	q=8	q=9	q=10
5	30.70	50.45	68.96	86.91	104.53	121.93	139.15	156.25	173.24
6	26.16	43.85	60.44	76.56	92.38	107.98	123.46	138.81	154.09
7	23.79	40.34	55.91	71.02	85.37	100.51	115.03	129.45	143.75
8	22.33	38.15	53.06	67.55	81.76	95.81	109.73	123.52	137.25
9	21.33	36.65	51.11	65.15	78.34	92.55	106.00	119.46	132.76
10	20.60	35.56	49.67	63.39	76.37	90.17	103.37	116.45	129.48
11	20.08	34.72	48.58	62.04	75.28	88.36	101.30	114.15	126.93
12	19.62	34.06	47.71	60.98	74.02	86.90	99.68	112.34	124.95
13	19.27	33.53	47.00	60.12	73.00	85.73	98.36	110.88	123.31
14	18.98	33.08	46.42	59.40	72.17	84.76	97.26	109.65	121.98
15	18.74	32.71	45.94	58.80	71.45	83.95	96.32	108.61	120.85
16	18.53	32.40	45.52	58.30	70.34	83.25	95.55	107.73	119.87
17	18.36	32.12	45.16	57.85	70.33	82.65	94.86	106.98	119.03
18	18.21	31.89	44.85	57.47	69.87	82.13	94.26	106.32	118.30
19	18.07	31.68	44.58	57.13	69.47	81.66	93.75	105.73	117.67
20	17.96	31.49	44.33	56.83	69.12	81.25	93.28	105.22	117.10
25	17.51	30.81	43.43	55.72	67.30	79.75	91.57	103.33	115.00
30	17.24	30.37	42.86	55.02	66.37	78.79	90.49	102.11	113.67

 $\alpha = 0.10 \quad p = 5$

n_o	q=2	q=3	q=4	q=5	q=6	q=7	q=8	q=9	q=10
6	45.25	74.90	102.86	130.09	156.89	183.39	209.65	235.79	261.72
7	38.56	65.12	90.21	114.66	138.70	162.50	186.15	209.58	232.90
8	34.95	59.77	83.27	106.16	128.70	150.99	173.12	195.06	216.94
9	32.69	56.37	78.81	100.70	122.23	143.54	164.69	185.70	206.61
10	31.12	54.00	75.70	96.88	117.72	138.36	158.82	179.16	199.37
11	29.97	52.24	73.40	94.03	114.37	134.47	154.42	174.28	194.00
12	29.08	50.88	71.62	91.85	111.78	131.47	151.03	170.52	189.84
13	28.39	49.31	70.20	90.09	109.70	129.10	148.33	167.49	186.53
14	27.82	48.93	69.04	88.66	108.01	127.14	146.14	165.04	183.78
15	27.35	48.21	68.07	87.48	106.60	125.51	144.31	162.94	181.54
16	26.95	47.59	67.26	86.47	105.42	124.14	142.75	161.24	179.62
17	26.62	47.07	66.57	85.62	104.40	122.98	141.42	159.75	177.97
18	26.32	46.61	65.96	84.87	103.50	121.95	140.27	158.45	176.56
19	26.07	46.21	65.44	84.22	102.74	121.06	139.23	157.32	175.30
20	25.85	45.85	64.97	83.65	102.05	120.28	138.36	156.32	174.20
25	25.03	44.59	63.28	81.56	99.57	117.41	135.10	152.67	170.19
30	24.52	43.78	62.22	80.23	98.01	115.60	133.04	150.39	167.69

TABLE VII (Continued)

 $\alpha = 0.10$ $p = 6$

n_o	q=2	q=3	q=4	q=5	q=6	q=7	q=8	q=9	q=10
7	62.54	104.12	143.53	181.97	219.37	257.40	294.64	331.76	368.58
8	53.42	90.76	126.20	160.82	194.34	228.72	262.28	295.59	328.81
9	48.40	83.29	110.44	148.84	180.79	212.41	243.93	275.04	306.13
10	45.18	78.42	110.07	141.01	171.49	201.73	231.72	261.56	291.26
11	42.92	74.99	105.56	135.44	164.91	194.13	223.14	251.93	280.69
12	41.23	72.42	102.16	131.26	159.38	188.42	216.65	244.70	272.69
13	39.94	70.43	99.54	128.00	156.11	183.93	211.57	239.10	266.48
14	38.91	68.83	97.42	125.39	153.31	180.35	207.54	234.56	261.46
15	38.06	67.51	95.67	123.24	150.46	177.39	204.16	230.78	257.33
16	37.36	66.42	94.23	121.46	148.31	174.95	201.33	227.67	253.85
17	36.76	65.49	93.00	119.93	146.51	172.84	199.02	225.05	250.97
18	36.26	64.59	91.93	118.60	144.95	171.03	196.97	222.74	248.44
19	35.82	64.00	91.01	117.47	143.59	169.49	195.17	220.73	246.20
20	35.43	63.39	90.21	116.47	142.40	168.11	193.61	219.00	244.29
25	34.06	61.22	87.30	112.86	138.11	163.13	187.98	212.70	237.33
30	33.21	59.97	85.49	110.63	135.45	160.06	184.51	208.78	233.04

 $\alpha = 0.05$ $p = 2$

n_o	q=2	q=3	q=4	q=5	q=6	q=7	q=8	q=9	q=10
3	12.18	18.70	24.55	30.09	35.45	40.68	45.81	50.87	55.86
4	10.70	16.65	22.00	27.07	31.37	36.75	41.45	46.07	50.64
5	9.97	15.63	20.73	25.57	30.23	34.79	39.26	43.67	48.02
6	9.53	15.02	19.97	24.66	29.19	33.61	37.95	42.22	46.45
7	9.24	14.62	19.46	24.05	28.49	32.83	37.08	41.26	45.40
8	9.04	14.33	19.10	23.62	27.99	32.26	36.44	40.57	44.64
9	8.88	14.11	18.83	23.30	27.62	31.84	35.99	40.09	44.08
10	8.76	13.94	18.61	23.05	27.33	31.51	35.61	39.65	43.64
11	8.67	13.81	18.44	22.85	27.10	31.25	35.32	39.33	43.29
12	8.59	13.70	18.30	22.68	26.90	31.03	35.03	39.07	43.00
13	8.52	13.60	18.19	22.54	26.75	30.85	34.87	38.84	42.76
14	8.47	13.53	18.10	22.42	26.61	30.70	34.71	38.66	42.56
15	8.42	13.46	18.01	22.33	26.50	30.57	34.57	38.50	42.38
16	8.38	13.40	17.94	22.24	26.40	30.45	34.43	38.36	42.23
17	8.35	13.35	17.87	22.17	26.31	30.35	34.32	38.24	42.10
18	8.32	13.30	17.82	22.10	26.23	30.27	34.23	38.13	41.99
19	8.28	13.26	17.77	22.04	26.16	30.19	34.14	38.04	41.88
20	8.26	13.23	17.72	21.98	26.10	30.12	34.07	37.95	41.79
25	8.17	13.10	17.55	21.79	25.97	29.86	33.78	37.63	41.44
30	8.11	13.01	17.44	21.65	25.72	29.69	33.59	37.42	41.21

TABLE VII (Continued)

 $\alpha = 0.05 \quad p = 3$

n_o	q=2	q=3	q=4	q=5	q=6	q=7	q=8	q=9	q=10
4	22.41	35.00	46.58	57.68	68.50	79.11	89.60	99.94	110.21
5	19.19	30.52	40.95	50.95	60.59	70.26	79.69	89.03	98.27
6	17.57	28.24	38.06	47.49	56.57	65.69	74.53	83.39	92.09
7	16.59	26.84	36.29	45.37	54.20	62.89	71.44	79.90	89.30
8	15.93	25.90	35.10	43.93	52.54	60.99	69.32	77.57	85.73
9	15.46	25.22	34.24	42.90	51.33	59.62	67.78	75.86	83.87
10	15.11	24.71	33.59	42.11	50.42	58.57	66.62	74.58	82.46
11	14.83	24.31	33.08	41.50	49.71	57.76	65.71	73.57	81.36
12	14.61	23.99	32.67	41.00	49.13	57.11	64.97	72.75	80.45
13	14.43	23.73	32.33	40.60	48.65	56.56	64.36	72.09	79.72
14	14.28	23.50	32.05	40.26	48.26	56.11	63.86	71.53	79.11
15	14.15	23.32	31.81	39.97	47.92	55.73	63.43	71.05	78.60
16	14.04	23.16	31.60	39.72	47.63	55.40	63.06	70.64	78.14
17	13.94	23.02	31.43	39.50	47.38	55.11	62.73	70.27	77.76
18	13.86	22.89	31.26	39.31	47.16	54.86	62.45	69.97	77.41
19	13.79	22.78	31.13	39.15	46.96	54.64	62.21	69.69	77.11
20	13.72	22.69	31.01	39.00	46.79	54.44	61.98	69.45	76.84
25	13.48	22.33	30.55	38.44	46.15	53.70	61.16	68.54	75.84
30	13.32	22.10	30.25	38.09	45.73	53.22	60.62	67.94	75.18

 $\alpha = 0.05 \quad p = 4$

n_o	q=2	q=3	q=4	q=5	q=6	q=7	q=8	q=9	q=10
5	35.39	56.10	75.36	93.97	112.17	130.11	147.81	165.39	182.80
6	30.06	48.62	65.90	82.60	98.93	115.03	130.94	146.69	162.34
7	27.31	44.69	60.89	76.56	91.88	106.98	121.90	136.71	151.39
8	25.61	42.24	57.77	72.77	87.46	101.94	116.23	130.43	144.50
9	24.45	40.57	55.62	70.17	84.42	98.46	112.32	126.08	139.74
10	23.62	39.34	54.04	68.26	82.19	95.90	109.46	122.91	136.24
11	22.98	38.41	52.84	66.81	80.48	93.95	107.27	120.46	133.57
12	22.48	37.67	51.90	65.66	79.14	92.41	105.54	118.55	131.45
13	22.08	37.08	51.13	64.73	78.04	91.15	104.12	116.98	129.74
14	21.75	36.59	50.50	63.95	77.13	90.12	102.97	115.69	128.32
15	21.47	36.17	49.97	63.30	76.37	89.26	101.93	114.59	127.14
16	21.24	35.82	49.51	62.76	75.73	88.51	101.14	113.67	126.10
17	21.03	35.52	49.12	62.28	75.16	87.87	100.42	112.87	125.22
18	20.86	35.26	48.78	61.86	74.68	87.31	99.80	112.17	124.46
19	20.70	35.02	48.47	61.50	74.25	86.82	99.25	111.56	123.79
20	20.56	34.82	48.21	61.17	73.87	86.38	98.75	111.02	123.18
25	20.06	34.06	47.23	59.98	72.47	84.78	96.95	109.01	120.49
30	19.74	33.59	46.61	59.21	71.58	83.74	95.79	107.71	119.57

TABLE VII (Continued)

 $\alpha = 0.05 \quad p = 5$

n_o	q=2	q=3	q=4	q=5	q=6	q=7	q=8	q=9	q=10
6	51.11	81.99	110.92	138.98	166.54	193.71	220.66	247.37	273.88
7	43.40	71.06	97.03	122.22	146.95	171.34	195.43	219.47	243.30
8	39.29	65.15	89.45	113.03	136.18	159.04	181.65	204.14	226.48
9	36.71	61.39	84.62	107.17	129.30	151.17	172.80	194.27	215.64
10	34.93	58.78	81.25	103.06	124.48	145.64	166.56	187.37	208.02
11	33.62	56.85	78.75	100.02	120.72	141.54	161.99	182.24	202.37
12	32.62	55.37	76.83	97.68	118.15	138.38	158.34	178.23	198.03
13	31.83	54.19	75.30	95.82	115.36	135.86	155.54	175.10	194.51
14	31.19	53.23	74.05	94.29	114.16	133.80	153.21	172.49	191.68
15	30.66	52.44	73.01	93.02	112.56	132.07	151.29	170.36	189.30
16	30.22	51.76	72.14	91.94	111.41	130.61	149.66	168.53	187.32
17	29.83	51.19	71.39	91.03	110.34	129.38	148.25	166.99	185.61
18	29.51	50.69	70.74	90.23	109.39	128.29	147.03	165.65	184.10
19	29.22	50.26	70.17	89.54	108.57	127.36	145.97	164.45	182.81
20	28.97	49.88	69.67	88.93	107.85	126.52	145.02	163.38	181.65
25	28.05	48.48	67.86	86.70	105.21	123.51	141.62	159.60	177.49
30	27.48	47.61	66.71	85.29	103.56	121.60	139.47	157.22	174.67

 $\alpha = 0.05 \quad p = 6$

n_o	q=2	q=3	q=4	q=5	q=6	q=7	q=8	q=9	q=10
7	69.62	112.59	153.28	192.75	231.50	269.98	308.03	345.81	383.45
8	59.24	97.92	134.42	169.94	204.36	239.37	273.52	307.61	341.44
9	53.56	89.72	123.37	157.11	189.33	222.14	254.13	286.00	317.69
10	49.95	84.42	117.03	148.77	179.37	210.85	241.48	271.84	302.13
11	47.43	80.69	112.18	142.85	173.10	202.85	232.46	261.80	291.03
12	45.56	77.90	108.56	138.40	167.79	196.84	225.65	254.25	282.76
13	44.11	75.74	105.72	134.94	163.72	192.15	220.37	248.39	276.24
14	42.96	74.00	103.46	132.19	160.44	188.39	216.09	243.67	271.04
15	42.03	72.59	101.61	129.90	157.76	185.31	212.63	239.72	266.75
16	41.24	71.41	100.07	128.01	155.52	182.73	209.70	236.48	263.15
17	40.58	70.40	98.76	126.38	153.60	180.51	207.23	233.73	260.07
18	40.02	69.54	97.63	125.01	151.36	178.63	205.09	231.36	257.48
19	39.53	68.80	96.64	123.80	150.55	176.99	203.21	229.26	255.19
20	39.11	68.13	95.77	122.73	149.29	175.53	201.60	227.45	253.19
25	37.58	65.79	92.69	118.93	144.78	170.35	195.69	220.92	246.00
30	36.63	64.33	90.77	116.57	141.17	167.13	192.04	218.82	241.49

TABLE VII (Continued)

 $\alpha = 0.01 \quad p = 2$

n_o	q=2	q=3	q=4	q=5	q=6	q=7	q=8	q=9	q=10
3	17.67	24.99	31.49	37.60	43.45	49.14	54.70	60.16	65.52
4	15.53	22.25	28.20	33.80	39.15	44.36	49.44	54.43	59.35
5	14.46	20.38	26.56	31.90	37.01	41.97	46.82	51.57	56.26
6	13.84	20.17	25.58	30.76	35.72	40.54	45.24	49.86	54.40
7	13.42	19.52	24.93	30.00	34.16	38.59	42.20	46.71	51.16
8	13.12	19.13	24.46	29.46	34.25	38.90	43.44	47.89	52.28
9	12.90	18.84	24.11	29.03	33.80	38.39	42.83	47.28	51.61
10	12.72	18.61	23.84	28.74	33.44	37.99	42.44	46.80	51.10
11	12.58	18.43	23.62	28.49	33.15	37.68	42.09	46.42	50.69
12	12.47	18.29	23.44	28.28	32.92	37.41	41.80	46.11	50.35
13	12.37	18.16	23.30	28.11	32.72	37.20	41.57	45.85	50.07
14	12.30	18.06	23.17	27.97	32.56	37.01	41.36	45.63	49.83
15	12.23	17.97	23.07	27.84	32.42	36.85	41.19	45.44	49.62
16	12.17	17.89	22.97	27.73	32.29	36.72	41.04	45.28	49.45
17	12.12	17.83	22.89	27.64	32.18	36.60	40.91	45.13	49.29
18	12.07	17.77	22.82	27.56	32.09	36.49	40.79	45.01	49.16
19	12.03	17.71	22.75	27.48	32.01	36.40	40.69	44.89	49.03
20	11.99	17.67	22.69	27.42	31.93	36.32	40.60	44.79	48.93
25	11.86	17.49	22.48	27.16	31.55	36.00	40.25	44.42	48.52
30	11.77	17.37	22.34	27.00	31.47	35.80	40.02	44.17	48.25

 $\alpha = 0.01 \quad p = 3$

n_o	q=2	q=3	q=4	q=5	q=6	q=7	q=8	q=9	q=10
4	30.17	44.00	56.58	68.55	80.14	91.49	102.60	113.57	124.41
5	25.75	38.23	49.56	60.36	70.80	81.01	91.03	100.91	110.68
6	23.53	35.30	46.00	56.17	66.02	75.65	85.10	94.42	103.61
7	22.20	33.53	43.83	53.62	63.11	72.38	81.47	90.45	99.30
8	21.31	32.34	42.37	51.91	61.14	70.17	79.03	87.76	96.38
9	20.67	31.48	41.32	50.67	59.73	69.58	77.25	85.82	94.27
10	20.19	30.84	40.52	49.73	58.65	67.37	75.91	84.34	92.69
11	19.82	30.33	39.90	49.01	57.81	66.42	74.86	83.20	91.43
12	19.52	29.93	39.40	48.42	57.14	65.66	74.03	82.28	90.42
13	19.28	29.50	39.00	47.93	56.58	65.03	73.33	81.51	89.59
14	19.07	29.12	38.65	47.53	56.13	64.51	72.75	80.87	88.90
15	18.90	28.80	38.36	47.18	55.73	64.07	72.27	80.34	88.31
16	18.75	28.58	38.11	46.89	55.39	63.69	71.83	79.86	87.79
17	18.62	28.41	37.89	46.63	55.10	63.36	71.47	79.47	87.36
18	18.51	28.25	37.70	46.41	54.84	63.07	71.14	79.12	86.97
19	18.41	28.12	37.54	46.21	54.61	62.80	70.87	78.80	86.64
20	18.32	28.00	37.39	46.04	54.41	62.58	70.60	78.52	86.33
25	17.99	27.85	36.83	45.38	53.55	61.74	69.55	77.48	85.20
30	17.79	27.56	36.47	44.96	53.16	61.18	69.05	76.81	84.48

TABLE VII (Continued)

 $\alpha = 0.01$ $p = 4$

n_0	$q=2$	$q=3$	$q=4$	$q=5$	$q=6$	$q=7$	$q=8$	$q=9$	$q=10$
5	45.48	67.99	88.52	108.33	127.50	146.51	165.13	183.54	201.82
6	38.41	58.52	77.05	94.81	112.11	129.07	145.79	162.32	178.70
7	34.80	53.66	71.05	87.72	103.35	119.86	135.53	151.06	166.42
8	32.58	50.86	67.33	83.30	98.86	114.12	129.14	143.99	158.73
9	31.08	48.61	64.78	80.27	95.36	110.17	124.74	139.17	153.43
10	30.00	47.12	62.92	78.07	92.92	107.28	121.54	135.60	149.57
11	29.18	45.99	61.50	76.38	90.97	105.07	119.07	132.89	146.62
12	28.54	45.10	60.39	75.06	89.33	103.34	117.13	130.77	144.28
13	28.03	44.38	59.48	73.98	88.09	101.92	115.56	129.03	142.39
14	27.60	43.79	58.74	73.09	87.05	100.75	114.26	127.60	140.81
15	27.24	43.29	58.12	72.34	86.19	99.78	113.16	126.40	139.49
16	26.94	42.86	57.59	71.71	85.45	98.94	112.24	125.35	138.36
17	26.68	42.50	57.13	71.17	84.82	98.23	111.42	124.46	137.41
18	26.46	42.18	56.72	70.68	84.26	97.59	110.71	123.70	136.56
19	26.26	41.90	56.38	70.26	83.79	97.04	110.11	123.02	135.81
20	26.08	41.65	56.06	69.89	83.36	96.55	109.56	122.42	135.13
25	25.44	40.74	54.92	68.52	81.77	94.75	107.55	120.19	132.73
30	25.03	40.17	54.19	67.64	80.75	93.60	106.26	118.78	131.15

 $\alpha = 0.01$ $p = 5$

n_0	$q=2$	$q=3$	$q=4$	$q=5$	$q=6$	$q=7$	$q=8$	$q=9$	$q=10$
6	63.59	96.63	127.33	156.94	185.34	214.26	242.31	270.07	297.64
7	53.60	83.24	110.79	137.35	163.28	188.83	214.03	238.93	263.64
8	48.35	76.11	101.91	126.76	151.07	174.97	198.58	221.92	245.07
9	45.09	71.61	96.28	120.07	143.30	166.13	188.68	211.02	233.21
10	42.86	68.51	92.37	115.39	137.88	159.97	181.82	203.43	224.84
11	41.23	66.22	89.50	111.93	133.85	155.44	176.71	197.81	218.68
12	39.99	64.47	87.29	109.29	130.76	151.90	172.78	193.43	213.96
13	39.00	63.08	85.53	107.18	128.32	149.13	169.68	190.01	210.18
14	38.21	61.95	84.10	105.46	126.31	146.86	167.08	187.15	207.06
15	37.55	61.01	82.90	104.02	124.66	144.93	164.97	184.83	204.48
16	37.00	60.23	81.90	102.82	123.26	143.34	163.17	182.85	202.34
17	36.52	59.55	81.04	101.78	122.04	141.98	161.57	181.15	200.45
18	36.12	59.37	80.30	100.88	120.99	140.77	160.31	179.65	198.86
19	35.77	58.46	79.65	100.10	120.08	139.76	159.14	178.35	197.42
20	35.46	58.01	79.09	99.41	119.29	138.82	158.11	177.22	196.18
25	34.33	56.38	77.00	96.91	116.36	135.49	154.38	173.09	191.64
30	33.62	55.37	75.70	95.34	114.52	133.40	152.02	170.48	188.82

TABLE VII (Continued)

 $\alpha = 0.01$ $p = 6$

n_o	$q=2$	$q=3$	$q=4$	$q=5$	$q=6$	$q=7$	$q=8$	$q=9$	$q=10$
7	84.51	130.23	173.00	214.41	254.35	294.86	334.36	373.45	412.35
8	71.34	112.45	150.89	188.07	224.50	260.43	295.88	331.06	365.98
9	64.27	102.74	138.71	173.53	207.62	241.23	274.41	307.34	340.08
10	59.81	96.51	130.87	164.10	196.59	228.79	260.53	291.97	323.20
11	56.71	92.15	125.34	157.47	188.32	219.93	250.64	280.98	311.19
12	54.43	88.92	121.24	152.51	183.14	213.36	243.13	272.79	302.24
13	52.68	86.41	118.04	148.67	178.64	208.24	237.44	266.46	295.22
14	51.29	84.41	115.48	145.56	175.06	204.06	232.85	261.33	289.58
15	50.16	82.77	113.38	143.04	172.11	200.75	229.06	257.12	284.98
16	49.22	81.41	111.65	140.92	169.51	197.92	225.90	253.59	281.09
17	48.41	80.26	110.16	139.13	167.53	195.49	223.20	250.62	277.86
18	47.74	79.27	108.91	137.60	165.73	193.46	220.87	248.03	275.04
19	47.15	78.40	107.80	136.26	164.17	191.64	218.95	245.80	272.61
20	46.63	77.65	106.83	135.09	162.80	190.07	217.07	243.87	270.47
25	44.80	74.37	103.36	130.88	157.87	184.44	210.76	236.82	262.71
30	43.67	73.30	101.21	128.26	154.80	180.93	206.73	232.45	257.88

SECTION VII

LIKELIHOOD RATIO TEST STATISTIC FOR MULTIPLE HOMOGENEITY OF THE COVARIANCE MATRICES

In this section, we discuss Pearson type approximations to the distribution of the likelihood ratio statistic for testing the hypothesis H_5 when $p_i = p(i = 1, \dots, q)$ and where H_5 was defined in Section II. This hypothesis is of interest in studying certain linear structures on the covariance matrices (see Krishnaiah and Lee [19]). The likelihood ratio statistic for testing H_5 is

$$v_5 = \frac{\prod_{i=1}^q \left| A_{ii}/n_i \right|^{n_i/2}}{\prod_{j=1}^k \left| \sum_{i=q_{j-1}+1}^{q_j} A_{ii}/n_j^* \right|^{n_j^*/2}} \quad (16)$$

where n_i and A_{ii} were defined in the preceding section, and

$$n_j^* = \sum_{i=q_{j-1}+1}^{q_j} n_i$$

The h-th moment of V_5 is given by

$$E(V_5^h) = \left[\frac{\prod_{\alpha=1}^k (n_{\alpha}^*)^{hn_{\alpha}^* p/2}}{\prod_{g=1}^q (n_{\alpha}^*)^{hn_g p/2}} \right] \left\{ \prod_{i=1}^p \prod_{\alpha=1}^k \left(\frac{\pi^{q_{\alpha}^*}}{\pi^{q_{\alpha-1}^*+1}} \cdot \frac{\Gamma[\frac{1}{2}(n_g + hn_g + 1 - i)]}{\Gamma[\frac{1}{2}(n_g + 1 - i)]} \cdot \frac{\Gamma[\frac{1}{2}(n_{\alpha}^* + 1 - i)]}{\Gamma[\frac{1}{2}(n_{\alpha}^* + hn_{\alpha}^* + 1 - i)]} \right) \right\} \quad (17)$$

As in the preceding section, the distribution of $V_5^{1/b}$ is approximated with the Pearson's Type I distribution where b is a properly chosen integer. In Tables VIII and IX, we have $n_i = n_0$ and $q = kd$. In Table VIII, α_1 is the value of α obtained if we use Pearson type approximation whereas α_2 is the value of α obtained by using Box's asymptotic series up to order n^{-13} , where

$$P \left[V_5^* \leq c_5 \mid H_5 \right] = (1-\alpha) \quad (18)$$

and

$$V_5^* = -2 \log V_5.$$

Table VIII indicates that the accuracy of the Pearson type approximation is sufficient for practical purposes. Using the above approximation, we computed tables for the values of c_5 where $\alpha = 0.05, 0.01, k = 2, 3, M = 1(1)20(5)30, M = n_0 - p, p = 1, 2, 3, 4$. These values are given in Table IX.

TABLE VIII
COMPARISON OF THE PEARSON TYPE APPROXIMATION WITH THE
ASYMPTOTIC EXPRESSION FOR THE DISTRIBUTION OF V_5^*

n ₀	q	k	p=1			p=2			p=3			p=4		
			c ₅	α ₁	α ₂	c ₅	α ₁	α ₂	c ₅	α ₁	α ₂	c ₅	α ₁	α ₂
10	6	2	8.20	0.05	0.0500	18.97	0.05	0.0500	34.63	0.05	0.0501	56.43	0.05	0.0503
10	6	2	11.89	0.01	0.0100	24.30	0.01	0.0100	41.77	0.01	0.0100	65.68	0.01	0.0101
10	6	3	9.90	0.05	0.0500	23.28	0.05	0.0500	42.79	0.05	0.0499	69.79	0.05	0.0498
10	6	3	13.85	0.01	0.0100	29.03	0.01	0.0100	50.53	0.01	0.0100	79.81	0.01	0.0099
20	6	2	8.01	0.05	0.0500	17.89	0.05	0.0500	31.45	0.05	0.0500	49.16	0.05	0.0500
20	6	2	11.62	0.01	0.0100	22.91	0.01	0.0100	37.93	0.01	0.0100	57.17	0.01	0.0100
20	6	3	9.70	0.05	0.0500	22.09	0.05	0.0500	39.30	0.05	0.0500	61.81	0.05	0.0499
20	6	3	13.57	0.01	0.0100	27.55	0.01	0.0100	46.39	0.01	0.0100	70.60	0.01	0.0100
30	6	2	7.94	0.05	0.0500	17.55	0.05	0.0500	30.54	0.05	0.0500	47.20	0.05	0.0500
30	6	2	11.53	0.01	0.0100	22.48	0.01	0.0100	36.81	0.01	0.0100	54.88	0.01	0.0100
30	6	3	9.63	0.05	0.0500	21.72	0.05	0.0500	38.28	0.05	0.0500	59.61	0.05	0.0500
30	6	3	13.47	0.01	0.0100	27.09	0.01	0.0100	45.17	0.01	0.0100	68.09	0.01	0.0100

TABLE IX
PERCENTAGE POINTS OF V_5^*

M	$\alpha = 0.05$			$\alpha = 0.01$		
	$p = 1$			$p = 1$		
	$k = 2$			$k = 2$		
	$q = 4$	$q = 6$	$q = 9$	$q = 4$	$q = 6$	$q = 9$
1	7.23	9.42	11.26	10.90	13.44	15.55
2	5.84	8.93	10.76	10.48	12.92	14.95
3	5.09	8.72	10.48	10.22	12.59	14.60
4	4.56	8.55	10.23	10.05	12.38	14.36
5	4.17	8.44	10.17	9.92	12.22	14.20
6	3.81	8.33	10.07	9.83	12.11	14.08
7	3.56	8.29	10.00	9.76	12.02	13.98
8	3.32	8.24	9.95	9.70	11.95	13.91
9	3.29	8.20	9.90	9.65	11.89	13.85
10	3.26	8.16	9.87	9.61	11.84	13.80
11	3.24	8.14	9.84	9.58	11.80	13.76
12	3.22	8.11	9.81	9.55	11.77	13.72
13	3.20	8.09	9.79	9.53	11.74	13.69
14	3.13	8.07	9.77	9.51	11.71	13.66
15	3.18	8.06	9.75	9.49	11.69	13.64
16	3.17	8.04	9.73	9.47	11.67	13.62
17	3.15	8.03	9.72	9.46	11.65	13.60
18	3.15	8.02	9.71	9.45	11.64	13.58
19	3.14	8.01	9.70	9.44	11.62	13.57
20	3.13	8.00	9.69	9.43	11.61	13.55
22	3.12	7.98	9.67	9.41	11.59	13.53
24	3.11	7.97	9.66	9.39	11.57	13.51
26	3.10	7.96	9.64	9.38	11.55	13.49
28	3.09	7.95	9.63	9.37	11.54	13.48
30	3.09	7.94	9.62	9.36	11.53	13.47

TABLE IX (Continued)

M	$\alpha = 0.05$			$\alpha = 0.01$		
	$p = 2$			$p = 2$		
	$k = 2$			$k = 2$		
	$q = 4$	$q = 6$	$q = 9$	$q = 4$	$q = 6$	$q = 9$
1	13.64	26.38	42.86	26.19	33.76	51.69
2	17.24	23.17	38.16	23.01	29.66	46.03
3	16.06	21.58	35.84	21.44	27.63	43.21
4	15.36	20.64	34.44	20.50	26.43	41.53
5	14.89	20.01	33.51	19.88	25.63	40.40
6	14.56	19.57	32.55	19.44	25.05	39.60
7	14.31	19.23	32.35	19.11	24.63	39.00
8	14.12	18.97	31.96	18.85	24.30	38.53
9	13.97	18.77	31.66	18.65	24.03	38.16
10	13.84	18.60	31.40	18.48	23.81	37.85
11	13.73	18.46	31.19	18.34	23.63	37.60
12	13.65	18.34	31.01	18.22	23.48	37.39
13	13.57	18.24	30.86	18.12	23.35	37.20
14	13.50	18.15	30.73	18.03	23.24	37.04
15	13.45	18.07	30.61	17.95	23.14	36.90
16	13.40	18.00	30.51	17.89	23.05	36.78
17	13.35	17.94	30.42	17.83	22.97	36.67
18	13.31	17.89	30.33	17.77	22.91	36.57
19	13.28	17.84	30.26	17.73	22.84	36.48
20	13.24	17.79	30.20	17.68	22.79	36.40
22	13.19	17.72	30.08	17.61	22.69	36.26
24	13.14	17.65	29.98	17.54	22.61	36.15
26	13.10	17.60	29.90	17.49	22.54	36.05
28	13.06	17.55	29.83	17.44	22.48	35.96
30	13.03	17.51	29.77	17.40	22.43	35.89

TABLE IX (Continued)

M	$\alpha = 0.05$			$\alpha = 0.01$		
	$p = 3$			$p = 3$		
	$k = 2$			$k = 2$		
	$q = 4$	$q = 6$	$q = 9$	$q = 4$	$q = 6$	$q = 9$
1	37.36	51.25	60.49	46.88	62.09	71.79
2	32.02	43.95	52.80	40.07	53.13	62.50
3	23.32	40.25	48.87	36.65	48.61	57.76
4	27.70	38.02	46.47	34.58	45.88	54.90
5	25.60	36.52	44.84	33.20	44.07	52.97
6	25.32	35.44	43.67	32.22	42.76	51.58
7	25.23	34.63	42.79	31.47	41.77	50.53
8	24.77	34.00	42.10	30.90	41.01	49.70
9	24.40	33.49	41.55	30.43	40.40	49.05
10	24.03	33.03	41.09	30.05	39.89	48.50
11	23.84	32.73	40.71	29.74	39.47	48.05
12	23.62	32.44	40.39	29.47	39.11	47.67
13	23.44	32.19	40.10	29.24	38.81	47.34
14	23.28	31.96	39.86	29.04	38.54	47.05
15	23.14	31.76	39.65	28.86	38.31	46.80
16	23.02	31.60	39.46	28.71	38.11	46.58
17	22.91	31.45	39.30	28.57	37.93	46.39
18	22.81	31.32	39.15	28.44	37.76	46.21
19	22.72	31.20	39.01	28.33	37.61	46.05
20	22.64	31.09	38.89	28.23	37.48	45.90
22	22.53	30.90	38.68	28.06	37.25	45.65
24	22.39	30.74	38.50	27.92	37.06	45.44
26	22.29	30.60	38.35	27.79	36.89	45.26
28	22.20	30.48	38.22	27.68	36.75	45.10
30	22.12	30.38	38.10	27.59	36.62	44.97

TABLE IX (Continued)

	P = 4			$\alpha = 0.05$			$\alpha = 0.01$					
	k = 2			k = 3			k = 2			k = 3		
	q = 4	q = 6	q = 9	q = 4	q = 6	q = 9	q = 4	q = 6	q = 9	q = 4	q = 6	q = 9
M												
1	50.43	54.13	99.17	140.36	73.06	98.57	114.18	157.80				
2	51.49	71.57	86.14	122.00	61.91	83.68	98.79	136.76				
3	46.80	55.17	79.22	112.27	56.16	75.97	90.72	125.72				
4	43.30	51.15	74.92	106.21	52.63	71.23	85.73	118.83				
5	41.93	50.41	71.95	102.04	50.24	68.00	82.29	114.11				
6	40.50	49.43	69.79	98.97	48.51	65.68	79.81	110.66				
7	39.41	54.32	68.15	96.63	47.19	63.91	77.91	108.04				
8	38.55	53.73	66.81	94.81	46.16	62.52	76.41	105.98				
9	37.48	52.78	65.80	93.31	45.33	61.39	75.20	104.30				
10	37.31	51.69	64.94	92.08	44.65	60.47	74.20	102.93				
11	36.53	51.33	64.21	91.06	44.08	59.70	73.36	101.77				
12	35.43	50.75	63.58	90.16	43.59	59.03	72.65	100.79				
13	36.08	50.28	63.04	89.42	43.17	58.47	72.03	99.94				
14	35.78	49.86	62.58	88.75	42.80	57.98	71.49	99.20				
15	35.51	49.49	62.17	88.16	42.48	57.55	71.02	98.53				
16	35.23	49.16	61.81	87.66	42.21	57.17	70.60	97.95				
17	35.07	48.87	61.43	87.20	41.95	56.82	70.23	97.44				
18	34.38	48.61	61.19	86.78	41.73	56.52	69.90	96.99				
19	34.71	48.37	60.92	86.40	41.52	56.25	69.60	96.56				
20	34.55	48.15	60.68	86.06	41.33	55.99	69.32	96.17				
22	34.23	47.78	60.26	85.47	41.01	55.56	68.85	95.53				
24	34.00	47.47	59.92	84.97	40.74	55.19	68.45	94.96				
25	33.87	47.20	59.61	84.55	40.51	54.88	68.09	94.48				
28	33.70	46.97	59.35	84.19	40.31	54.61	67.80	94.07				
30	33.55	46.76	59.13	83.87	40.14	54.37	67.54	93.70				

REFERENCES

1. Wilks, S.S., Certain Generalizations in the Analysis of Variance, Biometrika, Vol 24, pp 471-494, 1932.
2. Wilks, S.S., On the Independence of k sets of Normally Distributed Statistical Variables, Econometrica, Vol 3, pp 309-325, 1935.
3. Kendall, M.G. and Stuart, A., The Advanced Theory of Statistics, (third edition), Hafner Publishing Company, New York, 1947.
4. Johnson, N.L., Nixon, E., Amos, D.E. and Pearson, E.S., Table of Percentage Points of Pearson Curves, for Given $\sqrt{\beta_1}$ and β_2 , Expressed in Standard Measure, Biometrika, Vol 50, pp 459-497, 1963.
5. Box, G.E.P., A General Distribution Theory for a Class of Likelihood Criteria, Biometrika, Vol 36, pp 317-346, 1949.
6. Consul, P.C., The Exact Distributions of Likelihood Criteria for Different Hypotheses. In Multivariate Analysis - II (P.R. Krishnaiah, editor), Academic Press, 1969.
7. Mathai, A.M. and Rathie, P.N., The Problem of Testing Independence, Statistica, Vol 31, pp 673-688, 1971.
8. Davis, A.W. and Field, J.B.F., Tables of Some Multivariate Test Criteria, Tech. Report No. 32, Division of Mathematical Statistics, CSIRO, Australia, 1971.
9. Davis, A.W., Percentile Approximations for a Class of Likelihood Criteria, Biometrika, Vol 58, pp 349-356, 1971.
10. Lee, J.C., Krishnaiah, P.R. and Chang, T.C., Approximations to the Distributions of the Determinants of Real and Complex Multivariate Beta Matrices, ARL 75-, Aerospace Research Laboratoryies, Wright-Patterson AFB, Ohio, 1975.
11. Mauchly, J.W., Significant Test for Sphericity of a Normal n-Variate Distribution, Ann. Math. Statist., Vol 1, pp 204-209, 1940.
12. Mathai, A.M. and Rathie, P.N., The Exact Distribution for the Sphericity Test, J Statis. Res., (Dacca), Vol 4, pp 140-159, 1970.
13. Nagarsenker, B.N. and Pillai, K.C.S., The Distribution of the Sphericity Test Criterion, J. Multivariate Anal., Vol 3, pp 226-235, 1973.
14. Anderson, T. , An Introduction to Multivariate Statistical Analysis, Wiley, New York, 1958.
15. Korin, B.P., On the Distribution of a Statistic Used for Testing a Covariance Matrix, Biometrika, Vol 55, pp 171-178, 1968.

16. Nagarsenker, B.N. and Pillai, K.C.S., Distribution of the Likelihood Ratio Criterion for Testing a Hypothesis Specifying a Covariance Matrix, Biometrika, Vol 60, pp 359-364, 1973.
17. Korin, B.P., On Testing the Equality of k Covariance Matrices, Biometrika, Vol 56, pp 216-217, 1969.
18. Bishop, D.J., On a Comprehensive Test for the Homogeneity of Variances and Covariances in Multivariate Problems, Biometrika, Vol 31, pp 31-55, 1939.
19. Krishnaiah, P.R. and Lee, J.C., On Covariance Structures, Sankhyā (to appear), 1975.

APPENDIX

Let W ($0 \leq W \leq 1$) be a random variable whose moments are given by

$$E(W^h) = C \left[\prod_{j=1}^b y_j^{y_j} / \prod_{i=1}^a x_i^{x_i} \right] \frac{\prod_{i=1}^a \Gamma[X_i(1+h) + \xi_i]}{\prod_{j=1}^b \Gamma[Y_j(1+h) + \eta_j]}$$

where C is chosen such that $E(W^0) = 1$ and $\sum_{i=1}^a X_i = \sum_{j=1}^b Y_j$. Box (1949) gave an asymptotic expression for the distribution of $-2\rho \log W$, where ρ ($0 \leq \rho \leq 1$) is an arbitrary constant.

In this appendix we give asymptotic expression for the distribution of $-2\rho \log W$ with remaining term of order N^{-15} where ρ is chosen such that $\omega_1 = 0$, ω_r is given by

$$\omega_r = \frac{(-1)^{r+1}}{r(r+1)} \left\{ \sum_{i=1}^a \frac{B_{r+1}(\beta_i + \xi_i)}{(\rho x_i)^r} - \sum_{j=1}^b \frac{B_{r+1}(\epsilon_j + \eta_j)}{(\rho y_j)^r} \right\}$$

$\beta_i = (1 - \rho)X_i$, $\epsilon_j = (1 - \rho)Y_j$ and $B_r(h)$ is the Bernoulli polynomial of degree r and order unity.

Let $g_\nu(z)$ be the probability density function (p.d.f.) of the χ^2 distribution with ν degrees of freedom and N be the sample size. The p.d.f. of $-2\rho \log W$ is given by

$$l(z) = \sum_{r=0}^{14} S_r(z) + O(N^{-15})$$

where

$$S_0(z) = g_f(z), \quad S_1(z) = 0, \quad S_2(z) = \omega_2 [g_{f+4}(z) - g_f(z)]$$

$$S_3(z) = \omega_3 [g_{f+6}(z) - g_f(z)]$$

$$S_4(z) = \omega_4 [g_{f+8}(z) - g_f(z)] + \frac{1}{2} \omega_2^2 [g_{f+8}(z) - 2g_{f+4}(z) + g_f(z)]$$

$$S_5(z) = \omega_5 [g_{f+10}(z) - g_f(z)] + \omega_2 \omega_3 [g_{f+10}(z) - g_{f+6}(z) - g_{f+4}(z) + g_f(z)]$$

$$S_6(z) = \omega_6 [g_{f+12}(z) - g_f(z)] + \omega_4 \omega_2 [g_{f+12}(z) - g_{f+8}(z) - g_{f+4}(z) + g_f(z)]$$

$$+ \omega_3^2 \left[\frac{1}{2} g_{f+12}(z) - g_{f+6}(z) + \frac{1}{2} g_f(z) \right]$$

$$+ \omega_2^3 \left[\frac{1}{6} g_{f+12}(z) - \frac{1}{2} g_{f+8}(z) + \frac{1}{2} g_{f+4}(z) - \frac{1}{6} g_f(z) \right]$$

$$S_7(z) = \omega_7 [g_{f+14}(z) - g_f(z)] + \omega_5 \omega_2 [g_{f+14}(z) - g_{f+10}(z) - g_{f+6}(z) + g_f(z)]$$

$$+ \omega_4 \omega_3 [g_{f+14}(z) - g_{f+8}(z) - g_{f+6}(z) + g_f(z)]$$

$$+ \omega_3 \omega_2^2 \left[\frac{1}{2} g_{f+14}(z) - g_{f+10}(z) - \frac{1}{2} g_{f+8}(z) + \frac{1}{2} g_{f+6}(z) + g_{f+4}(z) - \frac{1}{2} g_f(z) \right]$$

$$S_8(z) = \omega_8 [g_{f+16}(z) - g_f(z)] + \omega_6 \omega_2 [g_{f+16}(z) - g_{f+12}(z) - g_{f+8}(z) + g_f(z)]$$

$$+ \omega_5 \omega_3 [g_{f+16}(z) - g_{f+10}(z) - g_{f+6}(z) + g_f(z)]$$

$$+ \omega_4^2 \left[\frac{1}{2} g_{f+16}(z) - g_{f+8}(z) + \frac{1}{2} g_f(z) \right]$$

$$+ \omega_4 \omega_2^2 \left[\frac{1}{2} g_{f+16}(z) - g_{f+12}(z) + g_{f+4}(z) - \frac{1}{2} g_f(z) \right]$$

$$+ \omega_3^2 \omega_2 \left[\frac{1}{2} g_{f+16}(z) - \frac{1}{2} g_{f+12}(z) - g_{f+10}(z) + g_{f+6}(z) \right.$$

$$\left. + \frac{1}{2} g_{f+4}(z) - \frac{1}{2} g_f(z) \right]$$

$$+ \omega_2^4 \left[\frac{1}{24} g_{f+16}(z) - \frac{1}{6} g_{f+12}(z) + \frac{1}{4} g_{f+8}(z) - \frac{1}{6} g_{f+4}(z) + \frac{1}{24} g_f(z) \right]$$

$$\begin{aligned}
S_9(z) = & \omega_4 [g_{f+18}(z) - g_f(z)] + \omega_7 \omega_2 [g_{f+18}(z) - g_{f+14}(z) - g_{f+4}(z) + g_f(z)] \\
& + \omega_6 \omega_3 [g_{f+18}(z) - g_{f+12}(z) - g_{f+6}(z) + g_f(z)] \\
& + \omega_5 \omega_4 [g_{f+18}(z) - g_{f+10}(z) - g_{f+8}(z) + g_f(z)] \\
& + \omega_5 \omega_2^2 \left[\frac{1}{2} g_{f+18}(z) - g_{f+14}(z) + \frac{1}{2} g_{f+10}(z) - \frac{1}{2} g_{f+8}(z) + g_{f+4}(z) - \frac{1}{2} g_f(z) \right] \\
& + \omega_4 \omega_3 \omega_2 [g_{f+18}(z) - g_{f+14}(z) - g_{f+12}(z) - g_{f+10}(z) + g_{f+8}(z) + g_{f+6}(z) \\
& \quad + g_{f+4}(z) - g_f(z)] \\
& + \omega_3^3 \left[\frac{1}{6} g_{f+18}(z) - \frac{1}{2} g_{f+12}(z) + \frac{1}{2} g_{f+6}(z) - \frac{1}{6} g_f(z) \right] \\
& + \omega_3 \omega_2^3 \left[\frac{1}{6} g_{f+18}(z) - \frac{1}{2} g_{f+14}(z) - \frac{1}{6} g_{f+12}(z) + \frac{1}{2} g_{f+10}(z) \right. \\
& \quad \left. + \frac{1}{2} g_{f+8}(z) - \frac{1}{6} g_{f+6}(z) - \frac{1}{2} g_{f+4}(z) + \frac{1}{6} g_f(z) \right]
\end{aligned}$$

$$\begin{aligned}
S_{10}(z) = & \omega_{10} [g_{f+20}(z) - g_f(z)] + \omega_8 \omega_2 [g_{f+20}(z) - g_{f+16}(z) - g_{f+4}(z) + g_f(z)] \\
& + \omega_7 \omega_3 [g_{f+20}(z) - g_{f+14}(z) - g_{f+6}(z) + g_f(z)] \\
& + \omega_6 \omega_4 [g_{f+20}(z) - g_{f+12}(z) - g_{f+8}(z) + g_f(z)] \\
& + \omega_6 \omega_2^2 \left[\frac{1}{2} g_{f+20}(z) - g_{f+16}(z) + \frac{1}{2} g_{f+12}(z) - \frac{1}{2} g_{f+8}(z) \right. \\
& \quad \left. + g_{f+4}(z) - \frac{1}{2} g_f(z) \right] \\
& + \omega_5^2 \left[\frac{1}{2} g_{f+20}(z) - g_{f+10}(z) + \frac{1}{2} g_f(z) \right] \\
& + \omega_5 \omega_3 \omega_2 [g_{f+20}(z) - g_{f+16}(z) - g_{f+14}(z) + g_{f+6}(z) + g_{f+4}(z) - g_f(z)] \\
& + \omega_4^2 \omega_2 \left[\frac{1}{2} g_{f+20}(z) - \frac{1}{2} g_{f+16}(z) - g_{f+12}(z) + g_{f+8}(z) + \frac{1}{2} g_{f+4}(z) - \frac{1}{2} g_f(z) \right]
\end{aligned}$$

$$\begin{aligned}
& + \omega_4 \omega_3^2 \left[\frac{1}{2} g_{f+20}(z) - g_{f+14}(z) - \frac{1}{2} g_{f+12}(z) + \frac{1}{2} g_{f+8}(z) + g_{f+4}(z) - \frac{1}{2} g_f(z) \right] \\
& + \omega_3^2 \omega_2^4 \left[\frac{1}{4} g_{f+20}(z) - \frac{1}{2} g_{f+16}(z) - \frac{1}{2} g_{f+14}(z) + \frac{1}{4} g_{f+12}(z) + g_{f+8}(z) \right. \\
& \quad \left. + \frac{1}{4} g_{f+4}(z) - \frac{1}{2} g_{f+6}(z) - \frac{1}{2} g_{f+4}(z) + \frac{1}{4} g_f(z) \right] \\
& + \omega_2^3 \omega_4 \left[\frac{1}{6} g_{f+20}(z) - \frac{1}{2} g_{f+16}(z) + \frac{1}{3} g_{f+12}(z) + \frac{1}{3} g_{f+8}(z) - \frac{1}{2} g_{f+4}(z) + \frac{1}{6} g_f(z) \right] \\
& + \omega_2^5 \left[\frac{1}{120} g_{f+20}(z) - \frac{1}{24} g_{f+16}(z) + \frac{1}{12} g_{f+12}(z) - \frac{1}{12} g_{f+8}(z) \right. \\
& \quad \left. + \frac{1}{24} g_{f+4}(z) - \frac{1}{120} g_f(z) \right]
\end{aligned}$$

$$\begin{aligned}
S_1(z) = & \omega_1 [g_{f+22}(z) - g_f(z)] + \omega_7 \omega_2 [g_{f+22}(z) - g_{f+18}(z) - g_{f+4}(z) + g_f(z)] \\
& + \omega_3 \omega_3 [g_{f+22}(z) - g_{f+16}(z) - g_{f+6}(z) + g_f(z)] \\
& + \omega_7 \omega_2^3 \left[\frac{1}{2} g_{f+22}(z) - g_{f+18}(z) + \frac{1}{2} g_{f+14}(z) - \frac{1}{2} g_{f+8}(z) + g_{f+4}(z) - \frac{1}{2} g_f(z) \right] \\
& + \omega_7 \omega_4 [g_{f+22}(z) - g_{f+14}(z) - g_{f+8}(z) + g_f(z)] \\
& + \omega_3 \omega_3^3 [g_{f+22}(z) - g_{f+12}(z) - g_{f+10}(z) + g_f(z)] \\
& + \omega_3 \omega_3 \omega_2 [g_{f+22}(z) - g_{f+18}(z) - g_{f+6}(z) + g_{f+12}(z) - g_{f+10}(z) \\
& \quad + g_{f+6}(z) + g_{f+4}(z) - g_f(z)] \\
& + \omega_3 \omega_4 \omega_2 [g_{f+22}(z) - g_{f+18}(z) - g_{f+14}(z) - g_{f+12}(z) + g_{f+10}(z) + g_{f+8}(z) + g_{f+4}(z) \\
& \quad - g_f(z)] \\
& + \omega_3 \omega_3^2 \left[\frac{1}{2} g_{f+22}(z) - g_{f+16}(z) - \frac{1}{2} g_{f+12}(z) + \frac{1}{2} g_{f+10}(z) \right. \\
& \quad \left. + g_{f+6}(z) - \frac{1}{2} g_f(z) \right]
\end{aligned}$$

$$\begin{aligned}
& + \omega_5 \omega_2^3 \left[\frac{1}{6} g_{f+22}(\delta) - \frac{1}{2} g_{f+18}(\delta) + \frac{1}{2} g_{f+14}(\delta) - \frac{1}{6} g_{f+12}(\delta) \right. \\
& \quad \left. - \frac{1}{6} g_{f+10}(\delta) + \frac{1}{2} g_{f+8}(\delta) - \frac{1}{2} g_{f+4}(\delta) + \frac{1}{6} g_f(\delta) \right] \\
& + \omega_4^2 \omega_3 \left[\frac{1}{2} g_{f+22}(\delta) - \frac{1}{2} g_{f+16}(\delta) - g_{f+14}(\delta) + g_{f+8}(\delta) + g_{f+6}(\delta) - \frac{1}{2} g_f(\delta) \right] \\
& + \omega_4 \omega_3 \omega_2^4 \left[\frac{1}{2} g_{f+22}(\delta) - g_{f+18}(\delta) - \frac{1}{2} g_{f+16}(\delta) + g_{f+12}(\delta) + g_{f+10}(\delta) - \frac{1}{2} g_{f+8}(\delta) \right. \\
& \quad \left. - g_{f+4}(\delta) + \frac{1}{2} g_f(\delta) \right] \\
& + \omega_3^2 \omega_2 \left[\frac{1}{6} g_{f+22}(\delta) - \frac{1}{6} g_{f+18}(\delta) - \frac{1}{2} g_{f+16}(\delta) + \frac{1}{2} g_{f+12}(\delta) \right. \\
& \quad \left. + \frac{1}{2} g_{f+10}(\delta) - \frac{1}{2} g_{f+6}(\delta) - \frac{1}{6} g_{f+4}(\delta) + \frac{1}{6} g_f(\delta) \right] \\
& + \omega_3 \omega_2^4 \left[\frac{1}{24} g_{f+22}(\delta) - \frac{1}{6} g_{f+18}(\delta) - \frac{1}{24} g_{f+16}(\delta) + \frac{1}{4} g_{f+14}(\delta) \right. \\
& \quad + \frac{1}{6} g_{f+12}(\delta) - \frac{1}{6} g_{f+10}(\delta) - \frac{1}{4} g_{f+8}(\delta) + \frac{1}{24} g_{f+6}(\delta) \\
& \quad \left. + \frac{1}{6} g_{f+4}(\delta) - \frac{1}{24} g_f(\delta) \right]
\end{aligned}$$

$$\begin{aligned}
S_{12}(\delta) = & \omega_{12} [g_{f+24}(\delta) - g_f(\delta)] + \omega_1 \omega_2 [g_{f+24}(\delta) - g_{f+20}(\delta) - g_{f+4}(\delta) + g_f(\delta)] \\
& + \omega_9 \omega_2 [g_{f+24}(\delta) - g_{f+18}(\delta) - g_{f+6}(\delta) + g_f(\delta)] + \omega_8 \omega_4 [g_{f+24}(\delta) - g_{f+16}(\delta) - g_{f+8}(\delta) + g_f(\delta)] \\
& + \omega_8 \omega_2^2 \left[\frac{1}{2} g_{f+24}(\delta) - g_{f+20}(\delta) + \frac{1}{2} g_{f+16}(\delta) - \frac{1}{2} g_{f+8}(\delta) + g_{f+4}(\delta) - \frac{1}{2} g_f(\delta) \right] \\
& + \omega_7 \omega_5 [g_{f+24}(\delta) - g_{f+14}(\delta) - g_{f+10}(\delta) + g_f(\delta)] \\
& + \omega_7 \omega_3 \omega_2 [g_{f+24}(\delta) - g_{f+20}(\delta) - g_{f+18}(\delta) + g_{f+14}(\delta) - g_{f+10}(\delta) + g_{f+6}(\delta) \\
& \quad + g_{f+4}(\delta) - g_f(\delta)]
\end{aligned}$$

$$\begin{aligned}
& + \omega_6^2 \left[\frac{1}{2} g_{f+24}(z) - g_{f+12}(z) + \frac{1}{2} g_f(z) \right] \\
& + \omega_6 \omega_4 \omega_2 [g_{f+24}(z) - g_{f+20}(z) - g_{f+16}(z) + g_{f+8}(z) + g_{f+4}(z) - g_f(z)] \\
& + \omega_6 \omega_2^3 \left[\frac{1}{2} g_{f+24}(z) - \frac{1}{2} g_{f+20}(z) + \frac{1}{2} g_{f+16}(z) - \frac{1}{3} g_{f+12}(z) + \frac{1}{2} g_{f+8}(z) \right. \\
& \quad \left. - \frac{1}{2} g_{f+4}(z) + \frac{1}{6} g_f(z) \right] \\
& + \omega_6 \omega_3^2 \left[\frac{1}{2} g_{f+24}(z) - g_{f+18}(z) + g_{f+6}(z) - \frac{1}{2} g_f(z) \right] \\
& + \omega_5^2 \omega_2 \left[\frac{1}{2} g_{f+24}(z) - \frac{1}{2} g_{f+20}(z) - g_{f+14}(z) + g_{f+10}(z) + \frac{1}{2} g_{f+4}(z) - \frac{1}{2} g_f(z) \right] \\
& + \omega_5 \omega_4 \omega_3 [g_{f+24}(z) - g_{f+18}(z) - g_{f+16}(z) - g_{f+14}(z) + g_{f+10}(z) + g_{f+8}(z) \\
& \quad + g_{f+6}(z) - g_f(z)] \\
& + \omega_5 \omega_3 \omega_2^2 \left[\frac{1}{2} g_{f+24}(z) - g_{f+20}(z) - \frac{1}{2} g_{f+18}(z) + \frac{1}{2} g_{f+12}(z) + \frac{1}{2} g_{f+10}(z) \right. \\
& \quad \left. + \frac{1}{2} g_{f+6}(z) + \frac{1}{2} g_{f+8}(z) - \frac{1}{2} g_{f+4}(z) - g_{f+2}(z) + \frac{1}{2} g_f(z) \right] \\
& + \omega_4^3 \left[\frac{1}{6} g_{f+24}(z) - \frac{1}{2} g_{f+16}(z) + \frac{1}{2} g_{f+8}(z) - \frac{1}{6} g_f(z) \right] \\
& + \omega_4^2 \omega_2^2 \left[\frac{1}{4} g_{f+24}(z) - \frac{1}{2} g_{f+20}(z) - \frac{1}{4} g_{f+16}(z) + g_{f+12}(z) - \frac{1}{4} g_{f+8}(z) \right. \\
& \quad \left. - \frac{1}{2} g_{f+4}(z) + \frac{1}{4} g_f(z) \right] \\
& + \omega_4 \omega_3 \omega_2 \left[\frac{1}{2} g_{f+24}(z) - \frac{1}{2} g_{f+20}(z) - g_{f+18}(z) - \frac{1}{2} g_{f+16}(z) + g_{f+14}(z) \right. \\
& \quad \left. + g_{f+12}(z) + g_{f+10}(z) - \frac{1}{2} g_{f+8}(z) - g_{f+6}(z) - \frac{1}{2} g_{f+4}(z) + \frac{1}{2} g_f(z) \right]
\end{aligned}$$

$$+ \omega_4 \omega_2^4 \left[\frac{1}{24} g_{f+24}(z) - \frac{1}{6} g_{f+20}(z) + \frac{5}{24} g_{f+16}(z) - \frac{5}{24} g_{f+12}(z) + \frac{1}{6} g_{f+8}(z) - \frac{1}{24} g_f(z) \right]$$

$$+ \omega_3^4 \left[\frac{1}{24} g_{f+24}(z) - \frac{1}{6} g_{f+18}(z) + \frac{1}{4} g_{f+12}(z) - \frac{1}{6} g_{f+6}(z) + \frac{1}{24} g_f(z) \right]$$

$$+ \omega_5^4 \omega_2^3 \left[\frac{1}{12} g_{f+24}(z) - \frac{1}{4} g_{f+20}(z) - \frac{1}{6} g_{f+18}(z) + \frac{1}{4} g_{f+16}(z) + \frac{1}{2} g_{f+14}(z) - \frac{1}{2} g_{f+10}(z) - \frac{1}{4} g_{f+8}(z) + \frac{1}{6} g_{f+6}(z) + \frac{1}{4} g_{f+4}(z) - \frac{1}{12} g_f(z) \right]$$

$$+ \omega_2^6 \left[\frac{1}{720} g_{f+24}(z) - \frac{1}{120} g_{f+20}(z) + \frac{1}{48} g_{f+16}(z) - \frac{1}{36} g_{f+12}(z) + \frac{1}{48} g_{f+8}(z) - \frac{1}{120} g_{f+4}(z) + \frac{1}{720} g_f(z) \right]$$

$$S_3(z) = \omega_3 [g_{f+6}(z) - g_f(z)] + \omega_4 \omega_2 [g_{f+26}(z) - g_{f+22}(z) - g_{f+18}(z) + g_f(z)]$$

$$+ \omega_3 \omega_2 [g_{f+26}(z) - g_{f+22}(z) - g_{f+18}(z) + g_f(z)]$$

$$+ \omega_4 \omega_2 [g_{f+26}(z) - g_{f+22}(z) - g_{f+18}(z) + g_f(z)]$$

$$+ \omega_4 \omega_2^2 \left[\frac{1}{2} g_{f+26}(z) - g_{f+22}(z) + \frac{1}{2} g_{f+18}(z) - \frac{1}{2} g_{f+14}(z) + g_{f+10}(z) - \frac{1}{2} g_f(z) \right]$$

$$+ \omega_3 \omega_2 [g_{f+26}(z) - g_{f+18}(z) - g_{f+14}(z) + g_f(z)]$$

$$+ \omega_3 \omega_2^2 [g_{f+26}(z) - g_{f+22}(z) - g_{f+18}(z) + g_{f+16}(z) - g_{f+10}(z) + g_{f+12}(z) + g_{f+4}(z) - g_f(z)]$$

$$+ \omega_4 \omega_2 [g_{f+26}(z) - g_{f+14}(z) - g_{f+12}(z) + g_f(z)]$$

$$+ \omega_7 \omega_4 \omega_2 [g_{f+26}(z) - g_{f+22}(z) - g_{f+18}(z) + g_{f+14}(z) - g_{f+12}(z) + g_{f+8}(z) + g_{f+4}(z) - g_f(z)]$$

$$+ \omega_1 \omega_3^2 \left[\frac{1}{2} g_{f+26}(z) - g_{f+22}(z) + \frac{1}{2} g_{f+14}(z) - \frac{1}{2} g_{f+12}(z) + g_{f+8}(z) - \frac{1}{2} g_f(z) \right]$$

$$+ \omega_7 \omega_2^3 \left[\frac{1}{6} g_{f+26}(z) - \frac{1}{2} g_{f+22}(z) + \frac{1}{2} g_{f+18}(z) - \frac{1}{6} g_{f+14}(z) - \frac{1}{6} g_{f+12}(z) \right. \\ \left. + \frac{1}{2} g_{f+8}(z) - \frac{1}{2} g_{f+4}(z) + \frac{1}{6} g_f(z) \right]$$

$$+ \omega_6 \omega_3 \omega_2 [g_{f+26}(z) - g_{f+22}(z) - g_{f+18}(z) - g_{f+14}(z) + g_{f+12}(z) + g_{f+10}(z) + g_{f+8}(z) - g_f(z)]$$

$$+ \omega_6 \omega_4 \omega_3 [g_{f+26}(z) - g_{f+22}(z) - g_{f+18}(z) - g_{f+14}(z) + g_{f+12}(z) + g_{f+8}(z) + g_{f+6}(z) - g_f(z)]$$

$$+ \omega_6 \omega_3 \omega_2^2 \left[\frac{1}{2} g_{f+26}(z) - g_{f+22}(z) - \frac{1}{2} g_{f+20}(z) + \frac{1}{2} g_{f+18}(z) + g_{f+16}(z) \right. \\ \left. - \frac{1}{2} g_{f+14}(z) - \frac{1}{2} g_{f+12}(z) + g_{f+10}(z) + \frac{1}{2} g_{f+8}(z) - \frac{1}{2} g_{f+6}(z) \right. \\ \left. - g_{f+4}(z) + \frac{1}{2} g_f(z) \right]$$

$$+ \omega_5 \omega_4^2 \left[\frac{1}{2} g_{f+26}(z) - g_{f+22}(z) - \frac{1}{2} g_{f+18}(z) + \frac{1}{2} g_{f+16}(z) + g_{f+14}(z) - \frac{1}{2} g_f(z) \right]$$

$$+ \omega_5 \omega_4 \omega_2^2 \left[\frac{1}{2} g_{f+26}(z) - g_{f+22}(z) - \frac{1}{2} g_{f+18}(z) + g_{f+16}(z) + g_{f+12}(z) - \frac{1}{2} g_{f+10}(z) \right. \\ \left. - g_{f+4}(z) + \frac{1}{2} g_f(z) \right]$$

$$+ \omega_5 \omega_3^2 \omega_2 \left[\frac{1}{2} g_{f+26}(z) - \frac{1}{2} g_{f+22}(z) - g_{f+20}(z) + \frac{1}{2} g_{f+18}(z) + \frac{1}{2} g_{f+14}(z) \right. \\ \left. + \frac{1}{2} g_{f+12}(z) + \frac{1}{2} g_{f+10}(z) - g_{f+6}(z) - \frac{1}{2} g_{f+4}(z) + \frac{1}{2} g_f(z) \right]$$

$$\begin{aligned}
& + \omega_3^2 \omega_3 \left[\frac{1}{2} g_{f+6}(z) - \frac{1}{2} g_{f+20}(z) - g_{f+16}(z) + g_{f+10}(z) + \frac{1}{2} g_{f+4}(z) - \frac{1}{2} g_f(z) \right] \\
& + \omega_3 \omega_2^4 \left[\frac{1}{24} g_{f+24}(z) - \frac{1}{6} g_{f+22}(z) + \frac{1}{4} g_{f+18}(z) - \frac{1}{24} g_{f+16}(z) - \frac{1}{6} g_{f+14}(z) \right. \\
& \quad \left. + \frac{1}{6} g_{f+12}(z) + \frac{1}{24} g_{f+10}(z) - \frac{1}{4} g_{f+8}(z) + \frac{1}{6} g_{f+4}(z) - \frac{1}{24} g_f(z) \right] \\
& + \omega_4 \omega_3^3 \left[\frac{1}{6} g_{f+24}(z) - \frac{1}{2} g_{f+20}(z) - \frac{1}{6} g_{f+18}(z) + \frac{1}{2} g_{f+14}(z) + \frac{1}{2} g_{f+12}(z) \right. \\
& \quad \left. - \frac{1}{6} g_{f+8}(z) - \frac{1}{2} g_{f+6}(z) + \frac{1}{6} g_f(z) \right] \\
& + \omega_4 \omega_3 \omega_2^3 \left[\frac{1}{6} g_{f+24}(z) - \frac{1}{2} g_{f+22}(z) - \frac{1}{6} g_{f+20}(z) + \frac{1}{3} g_{f+18}(z) + \frac{1}{2} g_{f+16}(z) \right. \\
& \quad \left. + \frac{1}{3} g_{f+14}(z) - \frac{1}{3} g_{f+12}(z) - \frac{1}{2} g_{f+10}(z) - \frac{1}{3} g_{f+8}(z) + \frac{1}{6} g_{f+6}(z) + \frac{1}{2} g_{f+4}(z) \right. \\
& \quad \left. - \frac{1}{6} g_f(z) \right] \\
& + \omega_4^2 \omega_3 \omega_2 \left[\frac{1}{2} g_{f+24}(z) - \frac{1}{2} g_{f+22}(z) - \frac{1}{2} g_{f+20}(z) - g_{f+18}(z) + \frac{1}{2} g_{f+16}(z) + g_{f+14}(z) \right. \\
& \quad \left. + g_{f+12}(z) + \frac{1}{2} g_{f+10}(z) - g_{f+8}(z) - \frac{1}{2} g_{f+6}(z) - \frac{1}{2} g_{f+4}(z) + \frac{1}{2} g_f(z) \right] \\
& + \omega_3^3 \omega_2^2 \left[\frac{1}{12} g_{f+24}(z) - \frac{1}{6} g_{f+22}(z) - \frac{1}{4} g_{f+20}(z) + \frac{1}{12} g_{f+18}(z) + \frac{1}{2} g_{f+16}(z) \right. \\
& \quad \left. + \frac{1}{4} g_{f+14}(z) - \frac{1}{4} g_{f+12}(z) - \frac{1}{2} g_{f+10}(z) - \frac{1}{12} g_{f+8}(z) + \frac{1}{4} g_{f+6}(z) \right. \\
& \quad \left. + \frac{1}{6} g_{f+4}(z) - \frac{1}{12} g_f(z) \right]
\end{aligned}$$

$$\begin{aligned}
& + \omega_3 \omega_2^5 \left[\frac{1}{120} g_{f+26}(z) - \frac{1}{24} g_{f+22}(z) - \frac{1}{120} g_{f+20}(z) + \frac{1}{12} g_{f+18}(z) \right. \\
& \quad + \frac{1}{24} g_{f+16}(z) - \frac{1}{12} g_{f+14}(z) - \frac{1}{12} g_{f+12}(z) + \frac{1}{24} g_{f+10}(z) + \frac{1}{12} g_{f+8}(z) \\
& \quad \left. - \frac{1}{120} g_{f+6}(z) - \frac{1}{24} g_{f+4}(z) + \frac{1}{120} g_f(z) \right]
\end{aligned}$$

$$S_4(z) = \omega_4 [g_{f+26}(z) - g_f(z)] + \omega_2 \omega_2 [g_{f+26}(z) - g_{f+24}(z) - g_{f+20}(z) + g_f(z)]$$

$$+ \omega_{11} \omega_3 [g_{f+28}(z) - g_{f+22}(z) - g_{f+16}(z) + g_f(z)]$$

$$+ \omega_{10} \omega_4 [g_{f+28}(z) - g_{f+26}(z) - g_{f+20}(z) + g_f(z)]$$

$$\begin{aligned}
& + \omega_{10} \omega_2^2 \left[\frac{1}{2} g_{f+28}(z) - g_{f+24}(z) + \frac{1}{2} g_{f+20}(z) - \frac{1}{2} g_{f+16}(z) + g_{f+12}(z) \right. \\
& \quad \left. - \frac{1}{2} g_f(z) \right]
\end{aligned}$$

$$+ \omega_9 \omega_5 [g_{f+28}(z) - g_{f+18}(z) - g_{f+10}(z) + g_f(z)]$$

$$\begin{aligned}
& + \omega_9 \omega_3 \omega_2 [g_{f+28}(z) - g_{f+24}(z) - g_{f+22}(z) + g_{f+18}(z) - g_{f+16}(z) + g_{f+14}(z) \\
& \quad + g_{f+12}(z) - g_f(z)]
\end{aligned}$$

$$+ \omega_8 \omega_6 [g_{f+28}(z) - g_{f+16}(z) - g_{f+12}(z) + g_f(z)]$$

$$+ \omega_8 \omega_3^2 \left[\frac{1}{2} g_{f+28}(z) - g_{f+22}(z) + \frac{1}{2} g_{f+16}(z) - \frac{1}{2} g_{f+12}(z) + g_{f+6}(z) - \frac{1}{2} g_f(z) \right]$$

$$+ \omega_8 \omega_6 \omega_2 [g_{f+28}(z) - g_{f+24}(z) - g_{f+20}(z) + g_{f+16}(z) - g_{f+12}(z) + g_{f+8}(z) + g_{f+4}(z) - g_f(z)]$$

$$\begin{aligned}
& + \omega_3 \omega_2^3 \left[\frac{1}{6} g_{f+28}(z) - \frac{1}{2} g_{f+24}(z) + \frac{1}{24} g_{f+20}(z) - \frac{1}{6} g_{f+16}(z) - \frac{1}{6} g_{f+12}(z) + \frac{1}{2} g_{f+8}(z) \right. \\
& \quad \left. - \frac{1}{2} g_{f+4}(z) + \frac{1}{6} g_f(z) \right] \\
& + \omega_7^2 \left[\frac{1}{2} g_{f+28}(z) - g_{f+14}(z) + \frac{1}{2} g_f(z) \right] \\
& + \omega_7 \omega_5 \omega_2 \left[g_{f+28}(z) - g_{f+24}(z) - g_{f+18}(z) + g_{f+10}(z) + g_{f+4}(z) - g_f(z) \right] \\
& + \omega_7 \omega_4 \omega_3 \left[g_f(z) - g_{f+28}(z) - g_{f+22}(z) - g_{f+26}(z) + g_{f+8}(z) + g_{f+6}(z) - g_f(z) \right] \\
& + \omega_7 \omega_3 \omega_2^2 \left[\frac{1}{2} g_{f+28}(z) - g_{f+24}(z) - \frac{1}{2} g_{f+22}(z) + \frac{1}{2} g_{f+20}(z) + g_{f+18}(z) \right. \\
& \quad \left. - g_{f+14}(z) + g_{f+10}(z) + \frac{1}{2} g_{f+8}(z) - \frac{1}{2} g_{f+6}(z) - g_{f+4}(z) + \frac{1}{2} g_f(z) \right] \\
& + \omega_6^2 \omega_2 \left[\frac{1}{2} g_{f+28}(z) - \frac{1}{2} g_{f+24}(z) - g_{f+16}(z) + g_{f+12}(z) + \frac{1}{2} g_{f+8}(z) - \frac{1}{2} g_f(z) \right] \\
& + \omega_6 \omega_3 \omega_3 \left[g_{f+28}(z) - g_{f+22}(z) - g_{f+18}(z) - g_{f+16}(z) + g_{f+12}(z) + g_{f+10}(z) + g_{f+6}(z) - g_f(z) \right] \\
& + \omega_6 \omega_4^2 \left[\frac{1}{2} g_{f+28}(z) - g_{f+26}(z) - \frac{1}{2} g_{f+16}(z) + \frac{1}{2} g_{f+12}(z) + g_{f+8}(z) - \frac{1}{2} g_f(z) \right] \\
& + \omega_6 \omega_4 \omega_2^2 \left[\frac{1}{2} g_{f+28}(z) - g_{f+24}(z) + \frac{1}{2} g_{f+16}(z) + \frac{1}{2} g_{f+12}(z) - g_{f+14}(z) + \frac{1}{2} g_f(z) \right] \\
& + \omega_6 \omega_3 \omega_2 \left[\frac{1}{2} g_{f+28}(z) - \frac{1}{2} g_{f+24}(z) - g_{f+22}(z) + g_{f+18}(z) + g_{f+10}(z) - g_{f+6}(z) \right. \\
& \quad \left. - \frac{1}{2} g_{f+4}(z) + \frac{1}{2} g_f(z) \right] \\
& + \omega_6 \omega_2^4 \left[\frac{1}{24} g_{f+28}(z) - \frac{1}{6} g_{f+24}(z) + \frac{1}{4} g_{f+20}(z) - \frac{5}{24} g_{f+16}(z) + \frac{5}{24} g_{f+12}(z) \right. \\
& \quad \left. - \frac{1}{4} g_{f+8}(z) + \frac{1}{6} g_{f+4}(z) - \frac{1}{24} g_f(z) \right]
\end{aligned}$$

$$\begin{aligned}
& + \omega_5^2 \omega_4 \left[\frac{1}{2} g_{f+28}(\beta) - \frac{1}{2} g_{f+24}(\beta) - g_{f+18}(\beta) + g_{f+10}(\beta) + \frac{1}{2} g_{f+8}(\beta) - \frac{1}{2} g_f(\beta) \right] \\
& + \omega_5^2 \omega_3 \left[\frac{1}{6} g_{f+28}(\beta) - \frac{1}{2} g_{f+24}(\beta) - \frac{1}{6} g_{f+18}(\beta) + \frac{1}{2} g_{f+16}(\beta) + \frac{1}{2} g_{f+12}(\beta) \right. \\
& \quad \left. - \frac{1}{6} g_{f+10}(\beta) - \frac{1}{2} g_{f+6}(\beta) + \frac{1}{6} g_f(\beta) \right] \\
& + \omega_5^2 \omega_2 \left[\frac{1}{4} g_{f+28}(\beta) - \frac{1}{2} g_{f+24}(\beta) + \frac{1}{4} g_{f+20}(\beta) - \frac{1}{2} g_{f+18}(\beta) + g_{f+14}(\beta) - \frac{1}{2} g_{f+10}(\beta) \right. \\
& \quad \left. + \frac{1}{4} g_{f+8}(\beta) - \frac{1}{2} g_{f+4}(\beta) + \frac{1}{4} g_f(\beta) \right] \\
& + \omega_5 \omega_4 \omega_3 \omega_2 \left[g_{f+28}(\beta) - g_{f+24}(\beta) - g_{f+22}(\beta) - g_{f+20}(\beta) + g_{f+16}(\beta) \right. \\
& \quad \left. + 2g_{f+14}(\beta) + g_{f+12}(\beta) - g_{f+8}(\beta) - g_{f+6}(\beta) - g_{f+4}(\beta) + g_f(\beta) \right] \\
& + \omega_5 \omega_3 \omega_2^2 \left[\frac{1}{6} g_{f+28}(\beta) - \frac{1}{2} g_{f+24}(\beta) - \frac{1}{6} g_{f+22}(\beta) + \frac{1}{2} g_{f+20}(\beta) \right. \\
& \quad \left. + \frac{1}{3} g_{f+18}(\beta) - \frac{1}{6} g_{f+16}(\beta) + \frac{1}{6} g_{f+12}(\beta) - \frac{1}{3} g_{f+10}(\beta) \right. \\
& \quad \left. - \frac{1}{2} g_{f+8}(\beta) + \frac{1}{6} g_{f+6}(\beta) + \frac{1}{2} g_{f+4}(\beta) - \frac{1}{6} g_f(\beta) \right] \\
& + \omega_4^3 \omega_2 \left[\frac{1}{6} g_{f+28}(\beta) - \frac{1}{6} g_{f+24}(\beta) - \frac{1}{2} g_{f+20}(\beta) + \frac{1}{2} g_{f+16}(\beta) + \frac{1}{2} g_{f+12}(\beta) \right. \\
& \quad \left. - \frac{1}{2} g_{f+8}(\beta) - \frac{1}{6} g_{f+4}(\beta) + \frac{1}{6} g_f(\beta) \right] \\
& + \omega_3^4 \omega_2 \left[\frac{1}{24} g_{f+28}(\beta) - \frac{1}{24} g_{f+24}(\beta) - \frac{1}{6} g_{f+22}(\beta) + \frac{1}{6} g_{f+18}(\beta) \right. \\
& \quad \left. + \frac{1}{4} g_{f+16}(\beta) - \frac{1}{4} g_{f+12}(\beta) - \frac{1}{6} g_{f+10}(\beta) + \frac{1}{6} g_{f+6}(\beta) \right. \\
& \quad \left. + \frac{1}{24} g_{f+4}(\beta) - \frac{1}{24} g_f(\beta) \right]
\end{aligned}$$

$$+ \omega_2^7 \left[\frac{1}{5040} g_{f+28}(z) - \frac{1}{720} g_{f+24}(z) + \frac{1}{240} g_{f+20}(z) \right. \\ \left. - \frac{1}{144} g_{f+16}(z) + \frac{1}{144} g_{f+12}(z) - \frac{1}{240} g_{f+8}(z) + \frac{1}{720} g_{f+4}(z) \right. \\ \left. - \frac{1}{5040} g_f(z) \right]$$

$$+ \omega_3^2 \omega_2^4 \left[\frac{1}{48} g_{f+28}(z) - \frac{1}{12} g_{f+24}(z) - \frac{1}{24} g_{f+20}(z) + \frac{1}{8} g_{f+16}(z) \right. \\ \left. + \frac{1}{6} g_{f+12}(z) - \frac{1}{16} g_{f+8}(z) - \frac{1}{4} g_{f+4}(z) - \frac{1}{16} g_f(z) \right. \\ \left. + \frac{1}{6} g_{f+16}(z) + \frac{1}{8} g_{f+8}(z) - \frac{1}{24} g_{f+4}(z) - \frac{1}{12} g_{f+2}(z) + \frac{1}{48} g_f(z) \right] \\ + \omega_4^2 \omega_3^2 \left[\frac{1}{4} g_{f+28}(z) - \frac{1}{2} g_{f+24}(z) - \frac{1}{2} g_{f+20}(z) + \frac{1}{4} g_{f+16}(z) + g_{f+12}(z) \right. \\ \left. + \frac{1}{4} g_{f+8}(z) - \frac{1}{2} g_{f+4}(z) - \frac{1}{2} g_{f+2}(z) + \frac{1}{4} g_f(z) \right]$$

$$+ \omega_4^2 \omega_2^3 \left[\frac{1}{12} g_{f+28}(z) - \frac{1}{4} g_{f+24}(z) + \frac{1}{12} g_{f+20}(z) + \frac{5}{12} g_{f+16}(z) \right. \\ \left. - \frac{5}{12} g_{f+12}(z) - \frac{1}{12} g_{f+8}(z) + \frac{1}{4} g_{f+4}(z) - \frac{1}{12} g_f(z) \right]$$

$$+ \omega_4^2 \omega_3^2 \omega_2^2 \left[\frac{1}{4} g_{f+28}(z) - \frac{1}{2} g_{f+24}(z) - \frac{1}{2} g_{f+20}(z) + g_{f+16}(z) \right. \\ \left. + \frac{3}{4} g_{f+12}(z) - \frac{3}{4} g_{f+8}(z) - g_{f+4}(z) + \frac{1}{2} g_{f+2}(z) \right. \\ \left. + \frac{1}{2} g_{f+4}(z) - \frac{1}{4} g_f(z) \right]$$

$$+ \omega_4^2 \omega_2^5 \left[\frac{1}{120} g_{f+28}(z) - \frac{1}{24} g_{f+24}(z) + \frac{3}{40} g_{f+20}(z) - \frac{1}{24} g_{f+16}(z) - \frac{1}{24} g_{f+12}(z) \right. \\ \left. + \frac{3}{40} g_{f+8}(z) - \frac{1}{24} g_{f+4}(z) + \frac{1}{120} g_f(z) \right]$$

$$\text{and } f = -2 \left\{ \sum_{i=1}^a \xi_i - \sum_{j=1}^b \eta_j - \frac{1}{2}(a-b) \right\}$$